

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active. Public Health (2006) 120, 27-32



MINI-SYMPOSIUM



Lessons learned from SARS: The experience of the Health Protection Agency, England

N.L. Goddard^a, V.C. Delpech^{a,*}, J.M. Watson^a, M. Regan^b, A. Nicoll^a

^aHealth Protection Agency Centre for Infections, 61 Colindale Avenue, London NW9 5EQ, UK ^bHealth Protection Agency North West, Vernon Pritchard Court, 57a Upper Northgate Street, Chester CH1 4EF, UK

Available online 16 November 2005



Summary The United Kingdom was assessed as a low risk country throughout the 2003 global SARS outbreaks. Despite this, 368 reports of potential SARS cases were made to the Health Protection Agency (HPA) between March and July 2003. The public health actions undertaken in response to these reports, the establishment of reporting mechanisms and the development of guidance documents were substantial. Lessons learned from mounting a UK response to SARS included: the importance of international collaboration; formation of a UK-wide, multidisciplinary Task Force; flexible case reporting mechanisms; integration of surveillance and laboratory data; generation of prompt and web-accessible guidance and advice; availability of surge capacity; and contingency planning. Lessons learned are being incorporated into the HPA's preparedness to prevent and control future newly emerging infectious disease threats.

© 2005 Published by Elsevier Ltd on behalf of The Royal Institute of Public Health.

Introduction

On 12 March 2003, the World Health Organisation (WHO) issued an unprecedented global alert regarding outbreaks of a severe pneumonia, subsequently characterised as Severe Acute Respiratory Syndrome (SARS), caused by the SARS-Coronavirus. SARS was the first new severe disease transmissible from person-to-person to emerge in the 21st Century. During the global outbreak, a total of 8096 probable cases of SARS and 774 deaths were reported from 29 countries and areas.¹ China (mainland), Taiwan and Hong Kong Special Administrative Region (SAR) experienced substantial outbreaks, although considerable numbers of cases were also reported from Canada, Singapore and Viet Nam.¹

The United Kingdom (UK) was assessed to be at low risk from SARS throughout the outbreak.² Between March and July 2003, 368 reports of suspected SARS cases were made to the Health Protection Agency (HPA).² Of these, only nine were initially classified as probable SARS cases, and only one patient, a 23-year-old male, was positive for SARS coronavirus (Co-V) on PCR testing and later showed evidence of seroconversion. Nonetheless,

0033-3506/\$ - see front matter © 2005 Published by Elsevier Ltd on behalf of The Royal Institute of Public Health. doi:10.1016/j.puhe.2005.10.003

^{*} Corresponding author. Tel.: +44 20 8327 6270; fax: +44 20 8200 7868.

E-mail address: valerie.delpech@hpa.org.uk (V.C. Delpech).

Guidance and advice

Provision of guidance and advice for healthcare professionals and the public formed an important component of the UK response to SARS during the outbreak period. The HPA website was the predominant mechanism for making guidance documents available in a timely manner.¹¹ Whilst the information and advice was generally well received, there are inherent difficulties in using a web site for disseminating information that is updated during an evolving situation. Users are required to undertake a proactive approach and check the website regularly for updates, which might prove particularly difficult in healthcare settings, where there is not regular access to the Internet within the workplace, or where there are regular changes of staff due to shift working. Furthermore, the constantly evolving situation highlighted the requirement to keep archives of web pages to create an audit trail of issued travel advisories and guidance documents.

There are also difficulties in maintaining an upto-date web resource when regular updates are being published on other external sites. For example, in England, travel guidance for the public was published by a number of different organisations; the HPA, the National Travel Health Network and Centre (NaTHNaC), the Department of Health and the Foreign and Commonwealth Office (FCO), as well by international organisations such as the WHO. Close collaboration was essential to ensure consistent messages were being conveyed.

In addition to information on the website, telephone queries from health care professionals and the media resulted in considerable demands on HPA staff time. During the peak period, approximately sixty telephone calls per day were received by the CFI which required public health advice; many more calls were made to the press office, and to regional and local centres. Management of the response process was challenging in ensuring that staffing was adequate and that all team members were kept up to date with current information. Regular team meetings at the beginning and end of each day were used to brief key staff and allocate tasks.

Surge capacity within the HPA

The Health Protection Agency was formed on 1 April 2003, almost one month after the WHO issued the global alert on 12 March 2003. All Divisions of the HPA were involved in the response to the first global outbreak. The level of response required was

considerable given the challenges to the resources of the new organisation, where areas of responsibility and lines of communication were still being developed.

Day-to-day operational response to the evolving outbreak involved epidemiologists who undertook strategic planning and response tasks in addition to their normal responsibilities; administrative support was largely provided on an ad hoc basis, which resulted in some lack of continuity. Particular areas of shortfall during the outbreak period included continuity of staff, especially experienced medical and non-medical epidemiologists, and lack of a centralised operations centre to facilitate sharing of epidemiological and virological information during the investigation of potential cases. These issues were addressed in the short term by utilising experienced staff undergoing higher medical training (e.g. Specialist Registrars), and by seconding a medical epidemiologist to the CFI's Respiratory Department.

CFI has responded to national and international infectious disease incidents for many years, primarily through the redeployment of staff working within other areas of the organisation, as outlined in its Major Outbreak and Incident Plan. The HPA Strategic Emergency Response Plan also makes provision to redeploy staff and operate shift working to provide round the clock cover.¹² Staff redeployment is likely to become easier to achieve politically as the level of escalation increases and the pressures become more widely acknowledged both within and outwith the organisation.

There is currently limited surge capacity to respond to an incident such as SARS that requires a large team over a prolonged period of time to prevent fatigue and potential burn-out of key staff involved in the response. Future re-emergence of SARS, or any other newly emergent infection would lead to the prompt escalation of surveillance activities, with corresponding demands on staffing and resources.

Contingency planning

The emergence of SARS illustrated the need to strike a balance between mounting a multi-agency outbreak response and ensuring NHS and partner organisations maintain continuity of key services. Similarities noted with the planning requirements for an Influenza pandemic and those of a phased escalation of response to SARS were recognised, developed and exercised.¹³ This set out specified actions at local, regional and national levels for the HPA at each level of escalation dependant upon

the extent of disease transmission abroad and in the UK. The HPA SARS Contingency Plan also complemented activities specified by the Department of Health for the NHS.¹⁴

Implementation of the phased contingency plan response raised a number of issues regarding NHS acute and primary care trust (PCT) preparedness. The most notable of these concerned the purchase and supply of personal protective equipment (PPE) consumable items such as masks, gowns and respirators, as clear lines of responsibility were difficult to identify. Implementation of infection control guidelines, including appropriate staff training were also highlighted as areas of concern.

Conclusions

SARS demonstrated the speed with which a readily transmissible disease could spread around the world during the 21st century, resulting in considerable social, economic and political impact in some countries. International collaboration was fundamental to the rapid identification of the causative agent, and also to the containment of SARS.

Despite the UK being assessed as low risk from SARS throughout the first global outbreak the public health response was substantial and provided many challenges. It has provided the opportunity to test many mechanisms already in place so that they will be further strengthened for the future. It has also prompted the drafting of detailed plans to respond to the re-emergence of SARS, or any other newly emergent infectious disease threat based on the lessons learned.

Both national and international collaboration proved vital in sharing timely information to inform the UK public health response. The use of electronic communication and teleconferencing was particularly effective in eliciting prompt responses from organisations and facilitating communication between expert groups (e.g. infection control and infectious disease experts) without the need to meet face-to face frequently. The model has been utilised subsequently for assessing the threat to the UK of avian influenza and is due to be adopted more formally as a component of the UK response to future threats.

Data from countries with substantial outbreaks demonstrated that basic public health and infection control measures such as contact tracing, infection control procedures, quarantine and voluntary home isolation were effective in controlling the outbreaks in the absence of a rapid diagnostic test, a vaccine or effective treatment. The outbreak highlighted that all levels of the healthcare system in the UK need to be prepared to respond; especially as the level of threat remains ever present in light of the continuing widespread avian influenza outbreaks in south east Asia, and the potential emergence of a strain of the influenza virus with pandemic potential.

Key points

1. The prompt sharing of epidemiological and virological data through international collaboration was fundamental to understanding this newly emergent infection and informing the UK public health response, and underpinned an early recognition of the importance of 'real time' clinical epidemiology and the need for rapid liaison between clinicians, virologists and public health colleagues.

2. The formation of a National (UK) SARS Task Force contributed to consistent approaches to professional and public information and advice across the UK, particularly on the use of case definitions and management algorithms.

3. Flexible case reporting mechanisms need to be implemented at central, regional and local level during an evolving outbreak to inform appropriate public health measures.

4. SARS demonstrated the need for an integrated system of surveillance information, laboratory data and local public health response data, including contract tracing, to enable coordination and strengthening of the response across the UK.

5. Dissemination of timely guidance and advice during an evolving outbreak is crucial and requires close collaboration of key organisations to ensure regular and consistent situation updates.

6. A prolonged level of response to an incident such as SARS requires adequate surge capacity. This is being addressed through the development of an HPA Strategic Emergency Response, and contingency plans.

7. The rapidly evolving understanding of the diagnosis, clinical management and transmission of SARS underlined the importance of having explicit planning assumptions upon which critical control points of escalation of response within contingency plans were based.

The ability to respond to any large outbreak in the UK requires substantial surge capacity to develop guidelines, establish robust reporting mechanisms, follow-up large numbers of contacts, respond to enquiries from health care professionals and the public, and to undertake risk assessment. The development of comprehensive contingency plans, clearly outlining the roles and responsibilities of key players has been undertaken at all levels within the UK, from the Health Departments through to NHS acute and primary care trusts in the light of SARS.

National surveillance, such as that undertaken by the HPA, is essential for monitoring the spread of an infection, however, the vigilance of primary health care professions is crucial in the early warning response. The UK was spared by not experiencing a substantial outbreak of SARS. Nonetheless, valuable lessons have been learnt which will ensure that it is better prepared in the event of future public health threats.

Acknowledgements

This paper was drafted with the support of the UK SARS Taskforce, whose comments were greatly appreciated. The UK SARS Taskforce membership includes representatives of: Health Departments: Department of Health (England), Department of Health, Social Services and Public Safety (Northern Ireland), National Assembly for Wales, Scottish Executive. National Centres: Communicable Disease Surveillance Centre (Northern Ireland), Communicable Disease Surveillance Centre (Wales), National Disease Surveillance Centre, Ireland, Scottish Centre for Infection and Environmental Health. Health Protection Agency (England): Business Division, Communicable Disease Surveillance Centre, Emergency Response Division, Local and Regional Services. Specialist & Reference Microbiology Division. Specialist Advisors: Clinical Infectious Disease, Infection Control. Communications: Department of Health, Health Protection Agency.

References

- World Health Organisation. Summary of probable SARS cases with onset of illness from 1 November 2002 to 31 July 2003 [Based on data as of the 31 December 2003]. Available at: http://www.who.int/csr/sars/country/table2004_04_21/ en/index.html [accessed 17 Nov 2004].
- Delpech V, Sedgwick JEC, Goddard NL, Simons P, Gopal R, Zambon MC, et al. Surveillance of severe acute respiratory syndrome (SARS) in a low-risk setting during the first global outbreaks. UK: Poster CFI, Health Protection Agency; Submitted for publication.
- SARS Expert Committee Report, Hong Kong government. Available at: http://www.sars-expertcom.gov.hk
- Van Loock F, Gill ON, Wallyn S, Nicoll A, Desenclos JC, Leinikki P. Roles and functions of a European Union Public Health Centre for communicable diseases and other threats to health. *Euro Surveill* 2002;7:78-84.
- World Health Organisation. *IHR revision proposals*; 12 January 2004. Available at: http://www.who.int/csr/ihr/ revisionprocess/working_paper/en/ [accessed 17 Nov 2004].
- Centers for Disease Control and Prevention, USA. Update: outbreak of severe acute respiratory syndrome—worldwide, 2003. MMWR Morb Mortal Wkly Rep 2003;52:241-48.
- 7. Chow Y, Anderson SR, Delpech V, Leese J, Horby P, Sedgwick JEC, et al. SARS: UK public health response past, present and future. *Commun Dis Public Health* 2003;6: 209-15.
- McDonald LC, Simor AE, Su IJ, Maloney S, Ofner M, Chen KT, et al. SARS in health care facilities, Toronto and Taiwan. *Emerg Infect Dis* 2004;10:777-81.
- 9. Gopalakrishna G, Choo P, Leo YS, Tay BK, Lim YT, Khan AS, et al. SARS transmission and hospital containment. *Emerg Infect Dis* 2004;10:395-400.
- Svoboda T, Henry B, Shulman L, Kennedy E, Rea E, Ng W, et al. Public health measures to control the spread of the severe acute respiratory syndrome during the outbreak in Toronto. N Engl J Med 2004;350:2352-61.
- Health Protection Agency website http://www.hpa.org.uk/ infections/topics_az/SARS.
- 12. Health Protection Agency. The HPA strategic emergency response plan. London: HPA; in press.
- Health Protection Agency. Interim contingency plan for severe acute respiratory syndrome (SARS); December 2003. Available at: http://www.hpa.org.uk/infections/topics_az/ SARS/pdfs/SARSContingencyDec03.pdf [accessed 18 Nov 2004].
- 14. Department of Health. Government Policy on SARS Available at: http://www.dh.gov.uk/PolicyAndGuidance/ HealthAndSocialCareTopics/SevereAcuteRespiratorySyndrome/SARSCONTENT_ID=4063833&chk=2F7%2Bq%2B.

Available online at www.sciencedirect.com SCIENCE DIRECT.