

Risk Reduction Framework for NHS Staff at risk of COVID-19 infection

Professor Kamlesh Khunti, Professor of Primary Care Diabetes & Vascular Medicine, University of Leicester *Email:* I&S

Dr Anne de Bono, President Faculty of Occupational Medicine, Consultant Occupational Physician. University Hospitals of Leicester *Email:*

Personal Data

Professor Ivan Browne, Honorary Professor in Public Health Practice, Director of Public Health and Sports Services, Leicester City Council, *Email:*

Irrelevant & Sensitive

Leslie Cove, Head of Equality and Inclusion – Advocacy and Activism, NHS England and NHS Improvement *Email:* leslie.cove@nhs.net

Andrew Foster, Executive Lead for Leadership and Interim Managing Director, NHS Leadership Academy *Email:* Andrew.foster2@leadershipacademy.nhs.uk

Professor Trish Greenhalgh, Nuffield Department of Primary Care Health Sciences, University of Oxford Radcliffe Observatory Quarter *Email:*

I&S

Professor Wasim Hanif, Professor Diabetes & Endocrinology, Consultant Physician, University Hospitals of Birmingham *Email:* Irrelevant & Sensitive@uhb.nhs.uk

Professor Azeem Majeed, Faculty of Medicine, School of Public Health. Chair - Primary Care and Public Health & Head of Department *Email:*

Irrelevant & Sensitive

Dr Chaand Nagpaul, General practitioner and Chair of the Council of the British Medical Association *Email:* CNagpaul@bma.org.uk

Professor NR, Chief Medical Officer and Consultant Cardiologist, University Hospitals Coventry & Warwickshire NHS Trust, Clifford Bridge Road, Coventry, CV2 2DX *Emails* NR@nhs.net NR@uhcw.nhs.uk

Mr Abdul Razaq, Interim Consultant in Public Health, Lancashire County Council *Email:* abdul.razaq@lancashire.gov.uk

Marcus Riddell, Head of Equality and Inclusion – Evidence and Policy, Acting Head of Temporary Staffing (to support COVID-19 tasks and staff bank development), NHS England and NHS Improvement *Email:* marcus.riddell@nhs.net

Professor Liam Smeeth, Professor of Clinical Epidemiology at London School of Hygiene and Tropical medicine. *Email:* I&S

Dr Sonya Wallbank, Head of Culture Transformation COVID 19 Clinical Health and Wellbeing, Lead, NHS England and NHS Improvement *Email:* sonya.wallbank@nhs.net

SARS -CoV-2 (COVID-19) is a new virus and its characteristics and effects are gradually being identified including risk factors for incidence and severity. WHO has classified COVID-19 as an international pandemic. The NHS is grateful for the contribution of its entire workforce at a time when it is confronted with the challenges of such a pandemic. Concerns have recently been raised in the UK due to disproportionately higher rates of COVID-19 in black and minority ethnic health populations (BAME) compared to white populations [1]. This has been particularly apparent amongst health care staff in the UK. The NHS is estimated to employ approximately 1.2-1.5 million staff, with BAME people making up 44% of the NHS workforce. Among all staff employed by the NHS, BAME account for approximately 21%, including approximately 20% among nursing and support staff and 44% among medical staff [2]. In a recent analysis of staff deaths, of the 106 included cases, 98 had patient facing roles, seven did not and this was unclear for one. In 89 cases, the individual had been working during the pandemic. The characteristics of the cases are presented in the Table[3]. The BAME health care staff affected were a heterogeneous group. Among the doctors, the specialties affected were surgery (five cases), general practice (four), emergency medicine and medicine (each two), and one each from histopathology, geriatrics, neurorehabilitation, paediatrics, and psychiatry. There were no anaesthetists or intensivists identified. Among the nursing staff, specialty was not always mentioned, but none were described as intensive care nurses. The relative ages, proportions of either sex and of ethnicity among the main groups of staff are shown in Table 1. Overall in view of the number and heterogeneity of the staff affected, it is difficult to infer further from the data.

Table. Age, gender and ethnicity of those who died from covid-19 among the main health and social care staff groups.

*For comparison, the approximate % of BAME the NHS workforce-[3] is included in the final row.

	Nurses and midwives	Healthcare support workers	Doctors and dentists	Other staff
Number	35	27	19	25
Age; yrs median (IQR [range])	51 (46-57 [23-70])	54 (42-64 [21-84])	62 (54-76 [36-79])	51 (34-58 [29-65])
Male; %	39	22	94	55
BAME; %	71	56	94	29
BAME workforce; %*	20	17	44	-

In addition, an emerging finding from systematic reviews and data from UK is indicating that particular comorbidities such as hypertension, cardiovascular disease and diabetes are more prevalent in people with severe COVID-19 [4, 5]. These comorbidities are also more prevalent in BAME populations and may explain the increased risk of morbidity and mortality in this group [1].

Public Health England has issued guidance which is updated regularly, to ensure implementation of measures to reduce the risk of community based and nosocomial contagion. These measures involve both behavioural instruction and the level of personal protective equipment (PPE) to be worn in different clinical environments. We first of all wish to ensure that the entire NHS workforce adheres to latest PHE recommendations to reduce the risk of acquiring COVID-19. PHE's original guidance also identified 3 factors used to guide managers in supporting conversations with staff about increased vulnerability; age >70year, selected underlying health conditions and pregnancy. Health conditions further divided into those 'extremely vulnerable' to COVID-19 for whom 'shielding' is currently required and those at 'increased risk of severe illness from COVID-19', requiring stringent social distancing (based on conditions previously identified as requiring an annual 'flu jab'). Ethnicity was not identified originally as a risk factor and is now to be included on basis of recent clinical evidence particularly ICU admissions data and healthcare staff fatalities. As far as we know evidence for ethnicity as an independent risk factor remains uncertain.

NHS employers guidance was issued recently for "extremely vulnerable" and "at risk" categories[6]. The guidance reiterated that preserving and protecting the health, safety and wellbeing of staff is critical for the NHS and to ensure they are mentally and physically healthy and protect their colleagues, patients and their families. The guidance has already made recommendations for staff who are at high risk of severe illness from COVID-19. Due to the early warning signs of a disparity in outcomes, there is a responsibility that the NHS has committed to address, which is to ensure the development of and consistent application of a risk reduction framework in order to guide employers as to how best protect a workforce and hence meet the needs of an equality impact assessment.[7]

In this paper we provide a COVID-19 Risk Reduction Framework for health care staff. The framework may help employers to complement risk assessment of their staff, particularly of high risk and vulnerable groups and ensure staff safety. *This Risk Reduction Framework needs to be used in conjunction with the NHS employers Guidance* and can be used as an aid to decision making and may be incorporated into existing risk templates which are already in use in many health care settings.

Method

A consensus group was set up and included a multidisciplinary group of primary and specialist care clinicians, members of equality and inclusions, NHS leadership, public health specialists, occupational health specialists, trust leads and researchers. In view of unavailability of robust data, the group made consensus recommendations based on the currently published literature and that could help to initiate further research and inform immediate public health and policy decisions to protect and save health care staff. The key aim was to develop a pragmatic simple Risk Reduction Framework that can be implemented in a health care setting (Figure). The Framework is based on the best available evidence which is limited, however, in view of the

serious and disproportionate impact of COVID-19, there is an urgent need to put in place immediate mitigations to try and prevent further serious consequences.

How to use the Risk Reduction Framework

Employers have a duty of care requiring them, as far as reasonably practicable, to secure the health safety and welfare of their employees. This includes an equitable approach to effective risk management and risk reduction of potential workplace hazards, for all staff which requires:

Workplace assessment of potential exposure to SARS-CoV-2 in the workplace, followed by application of an appropriate hierarchy of control measures, including elimination if possible, and reduction by hygiene measures, safe systems of work, selection and correct use of PPE, including necessary training and fit testing.

Workforce assessment to identify those individuals with potentially increased vulnerability to infection or poorer outcomes from COVID-19. Earlier guidance has already identified three categories of vulnerability, specific long term health conditions, older age and pregnancy. Recent evidence suggests that ethnicity, specifically a BAME background, may also be associated with increased vulnerability, particularly in the presence of other risk factors.[8]

Managers should seek to identify locally, in their team or service, those staff who may have increased vulnerability. Age and ethnicity of staff will already be known to managers but not necessarily any underlying health situation.

In normal circumstances it is not appropriate for managers to seek health information from staff beyond functional capabilities. However in the current situation health assessment by Occupational Health of all staff involved in this exercise will not be practicable and enquiry by managers of the presence of any underlying health condition would not be unreasonable, subject to caution about sensitivity and confidentiality.

It should not necessarily be assumed that staff, even with identified vulnerabilities, working in areas with the highest concentration of COVID-19 patients, such as ICU will be at the greatest risk. This depends upon the extent to which the risk of exposure is controlled by the measures above, including appropriate PPE.

Consideration of both aspects, workplace and individual, in supportive conversations between staff and managers which also take account of staff concerns and preferences should allow effective decision making about deployment, including adjustments in the workplace if appropriate. Some staff, including those who have underlying health conditions or particular concerns and anxieties about their health or work may require further assessment, advice and support in Occupational Health to facilitate decision making and risk reduction.

A Practical Route to Risk Reduction

The COVID-19 Risk Assessment Framework may help managers to make an assessment, with their staff of workplace and individual factors which should be considered as part of risk management and reduction. ***We recommend the assessment is done on a one to one consultation with their staff in a sensitive manner taking into consideration staff mental well-being. Employers need to ensure that cultural workplace factors are also taken into consideration so that staff have the confidence to raise their concerns.***

The framework complements existing guidance and takes account of emerging evidence that co-morbidities such as underlying health conditions and ethnicity may be associated with increased vulnerability to COVID-19. The framework assumes that community exposure is lower than in hospital settings, but this may not be the case, particularly if PPE is incorrectly used.

Key considerations to be taken account when making the assessment includes where the staff are based:

- Primary care or community Based?
- Hospital based ?
- In an environment where Aerosol Generating Procedures are performed?

Individual assessment

Four factors need to be considered:

1. Age

Those aged over 70 have already been identified by PHE as 'clinically vulnerable' and should take particular care to minimise contact with others outside their own household. Most will already be working remotely.

The Framework also identifies that risks are increased with age, and that this begins at an earlier age for those of BAME ethnicity.

2. Sex

Data globally and from UK are emerging that COVID-19 disproportionately affects men with UK data showing approximately 60% of people admitted to hospital being male.[9] **In addition these data also showed that female sex was associated with a 20% lower mortality.**

3. Underlying health conditions

'Clinically extremely vulnerable' People with health conditions in this group, identified by PHE, should have already received a letter about this or been contacted by their GP. They have been advised that shielding is required but the staff can still continue alternative duties from home. Some may be able to work from their shielded environment if practicable.

'Clinically vulnerable' people are those with underlying health conditions, or co-morbidities which place them at increased risk. Emerging evidence suggests that particular conditions, hypertension, cardiovascular disease, diabetes and chronic kidney disease are especially important risk factors, and these risk factors are increased in those of BAME ethnicity.[1] Obesity has now emerged as an independent risk factor for COVID-19 hospitalisation in the UK setting.[9]

4. Ethnicity

BAME ethnicity appears to be associated with increased risks, particularly in those with co-morbidities and at a younger age.

5. Pregnancy

Existing guidance identifies that pregnant women over 28 weeks should be regarded as at increased risk and recommended to stay at home.[10] For pregnant women with underlying health conditions at any stage of pregnancy a more precautionary approach is required and ethnicity should be included in the consideration and discussion between healthcare staff and managers.

Reducing workplace risk

The Risk Reduction Framework may help employers ensure appropriate adjustments are made to mitigate the risk of COVID-19 in high risk NHS staff. Employers will need to take into considerations local circumstances and other structural factors when making decisions in consultation with their staff. Some suggestions for high risk staff include a redeployment to lower risk environments. In the acute sector, as the NHS begins to focus on the development of green (COVID-19 negative clinical pathways) and blue (potential COVID-19 patient pathways) pathways, organisations will find that potential redeployment opportunities to green pathways will exist either within or across organisations and therefore where an organisation cannot redeploy staff within an organisation, it should consider redeployment across the health system. In primary care, roles which are not directly patient facing are emerging and could be used as redeployment opportunities, or measures such as avoiding direct contact in hot hubs might be appropriate precautions. In the community, higher levels of testing within nursing homes and community settings will identify areas of lower occupational risk. Managers may also consider referral to occupational health for assessment of risk factors such as blood pressure control and psychological support.

There is an urgent need for a greater level of understanding as to why workers, and indeed patients including those from a BAME background, appear to have a disproportionately increased morbidity and mortality from COVID-19 infection. A significant research and audit programme is also currently underway. This is the first attempt to create such a Risk Reduction Framework with currently available evidence and the Tool will be reviewed and updated in light of any evidence.

References

1. Khunti, K., et al., *Is ethnicity linked to incidence or outcomes of covid-19?* BMJ, 2020. **369**: p. m1548.
2. GOV.UK, *Ethnicity facts and figures: NHS Workforce*. . Available at <https://www.ethnicity-facts-figures.service.gov.uk/workforce-and-business/workforce-diversity/nhs-workforce/latest#by-ethnicity>. Accessed on 29th April 2020, 2020.
3. Cook, T., E. Kursumovic, and S. Lennane, *Exclusive: deaths of NHS staff from covid-19 analysed*. Available at <https://www.hsj.co.uk/exclusive-deaths-of-nhs-staff-from-covid-19-analysed/7027471.article>. Accessed on 29th April, 2020.
4. Docherty, A., et al., *Features of 16,749 hospitalised UK patients with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol*. Available at <https://www.hsj.co.uk/exclusive-deaths-of-nhs-staff-from-covid-19-analysed/7027471.article>. Accessed on 29th April 2020.
5. Yang, J., et al., *Prevalence of comorbidities and its effects in coronavirus disease 2019 patients: A systematic review and meta-analysis*. Int J Infect Dis, 2020. **94**: p. 91-95.
6. NHS Employers, *COVID-19 guidance for NHS workforce leaders*. Available at <https://www.nhsemployers.org/covid19>. Accessed on 1st May 2020.
7. Majeed A, Molokhia M, Pankhania P, Asanati K. Protecting the health of doctors during the Covid-19 pandemic. British Journal of General Practice 2020 (in press).
8. Platt L, Warwick R. Are some ethnic groups more vulnerable to COVID-19 than others? Institute for Fiscal Studies, Nuffield Foundation. 2020
9. Docherty AB, Harrison EM, Green CA et al. Features of 16,749 hospitalised UK patients with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol. <https://www.medrxiv.org/content/10.1101/2020.04.23.20076042v1>
- 10 <https://www.rcog.org.uk/>

Figure. See attached powerpoint slide