HEALTH SYSTEMS AND POLICY ANALYSIS

POLICY BRIEF 39

In the wake of the pandemic Preparing for Long COVID

Selina Rajan Kamlesh Khunti Nisreen Alwan Claire Steves Trish Greenhalgh Nathalie MacDermott Anna Sagan Martin McKee





INQ000249018_0001

Keywords: COVID-19 Health policy Health systems plans Public Health Symptom assessment Patient participation Research

Funding:

KK is supported by the National Institute for Health Research (NIHR) Applied Research Collaboration East Midlands (ARC EM) and the NIHR Leicester Biomedical Research Centre (BRC) in the United Kingdom.

Declarations of interest:

KK is Chair of the Ethnicity Subgroup of SAGE; KK and MM are members of Independent SAGE, all in the UK.

This policy brief is one of a new series to meet the needs of policy-makers and health system managers. The aim is to develop key messages to support evidence-informed policy-making and the editors will continue to strengthen the series by working with authors to improve the consideration given to policy options and implementation. © World Health Organization 2021 (acting as the host organization for, and secretariat of, the European Observatory on Health Systems and Policies)

Address requests about publications of the WHO Regional Office for Europe to:

Publications

WHO Regional Office for Europe UN City, Marmorvej 51 DK-2100 Copenhagen Ø, Denmark

Alternatively, complete an online request form for documentation, health information, or for permission to quote or translate, on the Regional Office web site (http://www.euro.who.int/pubrequest).

All rights reserved. The Regional Office for Europe of the World Health Organization welcomes requests for permission to reproduce or translate its publications, in part or in full.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either express or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use. The views expressed by authors, editors, or expert groups do not necessarily represent the decisions or the stated policy of the World Health Organization.

What is a Policy Brief?

A policy brief is a short publication specifically designed to provide policy makers with evidence on a policy question or priority. Policy briefs

- Bring together existing evidence and present it in an accessible format
- Use systematic methods and make these transparent so that users can have confidence in the material
- Tailor the way evidence is identified and synthesised to reflect the nature of the policy question and the evidence available
- Are underpinned by a formal and rigorous open peer review process to ensure the independence of the evidence presented.

Each brief has a one page key messages section; a two page executive summary giving a succinct overview of the findings; and a 20 page review setting out the evidence. The idea is to provide instant access to key information and additional detail for those involved in drafting, informing or advising on the policy issue.

Policy briefs provide evidence for policy-makers not policy advice. They do not seek to explain or advocate a policy position but to set out clearly what is known about it. They may outline the evidence on different prospective policy options and on implementation issues, but they do not promote a particular option or act as a manual for implementation.

In the wake of the pandemic: Preparing for Long COVID

Contents	page
Foreword	3
Key messages	5
Executive Summary	7
Policy Brief	9
1. Introduction	9
2. What do we know about Long COVID?	9
3. Policy and other responses in Europe	16
4. Implications for policy	20
References	21

Authors

Selina Rajan – European Observatory on Health Systems and Policies, London

Kamlesh Khunti – Diabetes Research Centre, University of Leicester, Leicester

Nisreen Alwan – University of Southampton, Southampton Claire Steves – King's College London, United Kingdom

Trish Greenhalgh – University of Oxford, Oxford

Nathalie MacDermott – King's College London

Alisha Morsella – Università Cattolica del Sacro Cuore, Rome Ester Angulo – Unidad de Investigación en Políticas y Servicios de Salud, Instituto Aragonés de Ciencias de la Salud (IACS) IIS Aragón Juliane Winkelmann – Department for Health Care Management, Technische Universität Berlin

Lucie Bryndová – Faculty of Social Sciences, Charles University, Prague

Inês Fronteira – Instituto de Higiene e Medicina Tropical, Lisbon Coralie Gandré – Institute for Research and Information in Health Economics (IRDES), Paris

Zeynep Or – Institute for Research and Information in Health Economics (IRDES), Paris

Sophie Gerkens – Belgian Health Care Knowledge Centre (KCE), Brussels

Anna Sagan – European Observatory on Health Systems and Policies, London

Jorge Simões – Instituto de Higiene e Medicina Tropical, Lisbon Walter Ricciardi – Università Cattolica del Sacro Cuore, Rome Antonio Giulio de Belvis – Università Cattolica del Sacro Cuore, Rome

Andrea Silenzi – Università Cattolica del Sacro Cuore, Rome, Italy Enrique Bernal-Delgado – Unidad de Investigación en Políticas y Servicios de Salud, Instituto Aragonés de Ciencias de la Salud (IACS) IIS Aragón

Francisco Estupiñán-Romero - Unidad de Investigación en Políticas y Servicios de Salud, Instituto Aragonés de Ciencias de la Salud (IACS) IIS Aragón

Martin McKee - European Observatory on Health Systems and Policies, London

Editor Anna Sagan

Series Editor Anna Sagan

Associate Editors

Josep Figueras Hans Kluge Suszy Lessof David McDaid Martin McKee Elias Mossialos Govin Permanand

Managing Editors

Jonathan North Lucie Jackson

The authors and editors are grateful to the reviewers who commented on this publication and contributed their expertise.

Print ISSN 1997-8065 Web ISSN 1997-8073

CORRIGENDUM

The following correction has been incorporated into the electronic file on 22nd March 2021:

• On page 11, the following sentence has been added: 'Further data is needed to establish the clinical relevance of such findings in children.'

Foreword

Over the past year the world has tracked the progress of the COVID-19 pandemic using data on cases and deaths. Yet we now know that these provide only a partial picture. Many people struggle to recover from the acute infection, suffering often disabling symptoms that last weeks or months and, in some cases, with disabilities that are likely to be very long lasting. Our understanding of this new condition, now termed Long COVID, is growing rapidly. For this, we owe a great deal to many people, but especially to those affected who have come together to document, analyse and report on the complex nature of this condition and its impact on their lives, as well as the health professionals, in some cases themselves suffering from Long COVID, who have initiated important research projects.

This policy brief seeks to raise awareness of Long COVID and to provide a resource for those in decision-making roles, setting out in basic terms what we know about this condition and what the policy options are for developing a response that leaves no one behind.

This is rapidly changing field and one where we have much to learn from the many initiatives being taken across Europe, some of which are described in this policy brief. We now need to implement and evaluate new models of care, building on the key messages from the 2018 Tallinn Conference that called on us to include, invest, and innovate and from the European Programme of Work 2020–25 United Action for Better Health. The legacy of COVID-19 will, sadly, be with us for a long time.

Hans Kluge Regional Director WHO Regional Office for Europe

Acknowledgments

We are grateful to Professor Trish Greenhalgh, University of Oxford, for her advice during the writing of this policy brief.

List of figures, tables and boxes

Tables

Table 1: Summary of selected studies on the prevalence of Long COVID in non-hospitalized patients	11
Table 2: Summary of selected studies on the prevalence of Long COVID in hospitalized patients postdischarge	12

Figures

Figure 1: The long-term effects of COVID-19	9
Boxes	
Box 1: What causes Long COVID?	10
Box 2: NICE guidelines on diagnosing Long COVID	16
Box 3: ICD-10 codes for use with Long COVID	16

Key messages

- COVID-19 can cause persistent ill-health. Around a quarter of people who have had the virus experience symptoms that continue for at least a month but one in 10 are still unwell after 12 weeks. This has been described by patient groups as "Long COVID".
- Our understanding of how to diagnose and manage Long COVID is still evolving but the condition can be very debilitating. It is associated with a range of overlapping symptoms including generalized chest and muscle pain, fatigue, shortness of breath, and cognitive dysfunction, and the mechanisms involved affect multiple system and include persisting inflammation, thrombosis, and autoimmunity. It can affect anyone, but women and health care workers seem to be at greater risk.
- Long COVID has a serious impact on people's ability to go back to work or have a social life. It affects their mental health and may have significant economic consequences for them, their families and for society.
- Policy responses need to take account of the complexity of Long COVID and how what is known about it is evolving rapidly. Areas to address include:
 - The need for multidisciplinary, multispecialty approaches to assessment and management;
 - Development, in association with patients and their families, of new care pathways and contextually appropriate guidelines for health professionals, especially in primary care to enable case management to be tailored to the manifestations of disease and involvement of different organ systems;
 - The creation of appropriate services, including rehabilitation and online support tools;
 - Action to tackle the wider consequences of Long COVID, including attention to employment rights, sick pay policies, and access to benefit and disability benefit packages;
 - Involving patients both to foster self-care and self-help and in shaping awareness of Long COVID and the service (and research) needs it generates; and
 - Implementing well-functioning patient registers and other surveillance systems; creating cohorts of patients; and following up those affected as a means to support the research which is so critical to understanding and treating Long COVID.

Executive summary

Long COVID is an emerging condition that is not yet well understood but that can be severely disabling

A proportion of people experience persisting ill health following the acute manifestations of COVID-19. There may be several reasons for this, but among them is the condition that is now termed Long COVID. The mechanisms by which the infection leads to Long COVID are also not yet fully understood. There is growing evidence that the virus can cause direct organ damage but also give rise to an abnormal response, increasing blood clotting and release of inflammatory substances. This can affect many different body systems, in particular the heart, lungs and brain.

Long COVID can be severely disabling, and those suffering from it report functional disabilities. Although there is no simple symptom or test for diagnosing it, many people experience severe fatigue and a range of troubling physical symptoms that make it difficult for those who are employed to return to work. This has obvious economic consequences.

A significant number of adults with COVID-19 suffer from Long COVID but exact numbers are currently difficult to estimate

Estimates vary as to how common Long COVID is, but about a quarter of those with COVID-19 have continuing symptoms 4–5 weeks after testing positive, with about one in 10 experiencing symptoms after 12 weeks. The reasons why some people get Long COVID and others do not remain unknown, although it is associated with increasing age, the number of symptoms in the acute phase, and being female, among other characteristics. Children can also, rarely, be affected by a separate multisystem condition. Existing surveillance systems developed to track the course of the pandemic have, so far, generally not recorded data on this condition, in part because of the lack of an agreed case definition.

An emerging consensus on best practice points to the importance of multidisciplinary and multispecialty approach to assessment and management

Existing guidelines propose that the initial management of Long COVID should be in primary care and should include a series of investigations both to characterize how the individual is affected and to exclude other conditions that may coexist. This should be universally accessible. Management should then be tailored to the manifestations of disease, including investigation and referral for signs of involvement of different organ systems. Particular attention must be paid to actions aimed at narrowing inequalities in Long COVID, including employment rights and sick pay policies, particularly for health care workers who are more at risk. At present very few countries have specific employment benefit packages to support patients to access disability benefits.

Different European countries have been responding to the emergence of this new condition via organized health systems responses

This policy brief lists a number of responses that have been adopted in different European countries. These include development of new guidelines for assessing and managing Long COVID, development of new care pathways, as well as creation of post-COVID clinics and online support tools. There are also examples of initiatives undertaken by local service providers and professional bodies that other countries can learn from.

Patients suffering from Long COVID have been instrumental in shaping awareness, research and service provision

There is a growing recognition of the importance of engagement with those affected, with a number of patient communities operating online, as well as groups that bring together health workers who have been affected and who have played an important role in generating and exchanging knowledge. Self-help groups have been created in many countries and have helped to articulate the patient voice and provide support.

Further research remains a high priority

Given the many unanswered questions about this condition, research is a high priority. The policy brief describes some of the ongoing studies being undertaken in Europe, following up those affected and conducting detailed clinical assessments. Some of these are creating cohorts of patients, something that will be of great value for the evaluation of potential treatments.

POLICY BRIEF

1. Introduction

At least 1 in 5 people infected with COVID-19 experience persisting ill health following the acute phase of infection (Office for National Statistics, 2020). There may be several reasons for this, but among them is the condition that is now termed Long COVID. As the scale of disease burden attributable to Long COVID becomes recognized, health systems in many countries are seeking to understand the implications in the medium and long term and are asking how they should respond. This policy brief, written for decision-makers, is in two parts. First, it summarizes what is known about this condition, reviewing evidence on the mechanisms involved, who and what proportion of those infected get it, how it affects them, and what is known about diagnosis and treatment. Second, it brings together examples of how European countries are responding to this new condition.

This policy brief is not a substitute for either the detailed guidelines for clinicians or the resources for patients that have been developed in several countries, some of which are referred to later in this brief. As this is a rapidly developing topic, this brief offers a primer for those audiences who are involved in planning a policy response to this emerging condition.

2. What do we know about Long COVID?

2.1. Causes and definition

Mechanisms by which Long COVID develops have not yet been fully understood but some of their harmful consequences are recognized

There is mounting evidence that Long COVID is both common and debilitating (as outlined in sections 2.3 and 2.4 and summarized in Fig. 1). Its development and severity do not seem to correlate with the extent and nature of symptoms during the acute phase of the infection (Greenhalgh et al., 2020), although it is more common in hospitalized patients (Sudre et al., 2020). The exact mechanisms by which Long COVID develops remain unknown (see Box 1).

Symptoms include generalized pain, fatigue, shortness of breath, chest pain, muscle aches, palpitations, persisting high temperature and cognitive dysfunction, among others (Davis et al., 2020; Greenhalgh et al., 2020; Office for National Statistics, 2020), clustering and overlapping, often changing over time, and with evidence of involvement of almost any system within the body. The unpredictable and debilitating nature of this clinical picture can also lead to mood disorders. The worrying aspects of Long COVID include the multisystem pathology (National Institute for



Fig. 1 The long-term effects of COVID-19

Source: Reproduced with permission from SIMPLECOVID (2020)

Health and Clinical Excellence, 2020) that is associated with a variety of harmful consequences including scarring in multiple organs (especially the lungs, heart, kidneys, liver, adrenal glands and gastrointestinal tract) (Dennis et al., 2020), development of autoimmunity (British Society for Immunology, 2020) and blood clots causing strokes or other tissue damage (Libby & Lüscher, 2020).

Box 1: What causes Long COVID?

Available evidence shows that SARS-CoV-2 can trigger a hypercoagulable state and endothelial dysfunction (Libby & Lüscher, 2020), resulting in widespread formation of thrombi, and dysregulation of the immune response leading to a so-called cytokine storm in hospitalized patients and dysregulated inflammation in milder cases (British Society for Immunology, 2020; Gupta et al., 2020). In essence, the body produces pro-inflammatory cytokines, which are designed to kill the virus but are also unintentionally harmful, especially when not counter-balanced by the production of anti-inflammatory cytokines. There are many precedents of acute viral infections leading to long-term sequelae through immune and inflammatory responses, such as measles and subacute sclerosing panencephalitis (Bellini et al., 2005). Others include Ebola virus, Chikungunya virus and Epstein-Barr virus, which itself has been proposed as a potential trigger for an autoimmune reaction that leads to multiple sclerosis (Altmann & Boyton, 2020). However, the consequences of infection with SARS-CoV-2 are especially widespread, with many, and particularly the cardiovascular complications, seeming to relate to the widespread presence of angiotensin-converting enzyme 2 (ACE-2) receptors, which are highly expressed in the heart and lungs and act as the binding site for coronaviruses, including SARS-CoV and SARS-CoV-2, increasing the risk of endothelial damage and thrombo-inflammation (British Society for Immunology, 2020; Gupta et al., 2020).

There is no internationally agreed definition of Long COVID

In early literature a variety of terms are used, including "post-COVID syndrome". The National Institute for Health and Clinical Excellence (NICE), the Scottish Intercollegiate Guidelines Network, and the Royal College of General Practitioners include two conditions within the umbrella of Long COVID (Ongoing symptomatic COVID-19: signs and symptoms of COVID-19 from 4 to 12 weeks; and Post-COVID-19 syndrome: signs and symptoms that develop during or after an infection consistent with COVID-19, continue for more than 12 weeks and are not explained by an alternative diagnosis). The French Haute Autorité de Santé specifies three criteria to identify patients suffering from "symptômes prolongés de la Covid-19": having presented with a symptomatic form of Covid-19; presenting with one or more initial symptoms 4 weeks after the start of the disease; and none of these symptoms can be explained by another diagnosis (Haute Autorité de Santé, 2021). However, there is no internationally agreed definition and this syndrome, however it is termed, is commonly associated with the persistence of a range of heterogeneous symptoms, some of which may be due to other problems, and the choice of 12 weeks is arbitrary (Alwan & Johnson, 2020).

2.2. Who is more susceptible to Long COVID?

Susceptibility in adults appears to increase with age, the number of symptoms in the acute phase, body mass index and being female

Developing symptomatic Long COVID is not dependent on the severity of the initial COVID-19 infection or the duration of symptoms associated with it. A German study found that 78% of people known to have had COVID-19 had evidence of myocardial involvement based on a detailed assessment, and two thirds of these had not required hospital admission. There was no correlation with the severity of the initial infection (Puntmann et al., 2020). However, not everyone seems to be equally susceptible. The COVID Symptom Study, based at King's College London, which has recruited 4 million people who report COVID-19 symptoms and test results on an app, found that the strongest predictors of persistent symptoms were increasing age, followed by the number of symptoms in the acute phase and, particularly, hoarseness of voice, greater body mass index, shortness of breath and being female (although men are more likely to die in the acute illness). Individuals with asthma were also reported to be at greater risk of developing Long COVID compared with those without. Among those over 70, the leading risk factors were loss of smell, early fever, hoarseness of voice and comorbidities, particularly those involving the heart and lung (Sudre et al., 2020). A study from the United States, based on telephone interviews with 292 people who had tested positive for COVID-19 but who were never admitted to hospital, found that 35% had not returned to their usual state of health after a median of 16 days from their original test. This increased steadily from 26% among those aged 18-34 years, to 32% among those aged 35-49 years, and 47% among those aged ≥50 years (Tenforde et al., 2020). Other surveys also suggest that increasing age, previous health status, presence of comorbidities and the number of symptoms during the acute phase, are all associated with the number of Long COVID symptoms an individual may subsequently develop (Goërtz et al., 2020).

2.3. How common is Long COVID?

Estimating prevalence of Long COVID is not easy due to differences in methods used in surveys

There are large variations in the symptoms and reported prevalence in studies of Long COVID because a common definition is still lacking (see section 2.1) and because surveys include people recruited in different ways, and record different sets of symptoms reported at different follow-up periods, undergoing different types of investigation, with many reports based on small samples that may not be particularly representative of those affected. Some survey participants have undergone especially detailed phenotyping including biomarker and radiological analyses, and it may be difficult to know whether any abnormalities detected are chance findings or are associated with their symptoms and whether these will resolve with time or persist. A living systematic review of 28 studies, 61% of which were from Europe and including 9 442 adults with COVID-19, noted that while Long COVID seemed to affect both previously hospitalized and nonhospitalized populations, the quality of evidence was low, with a high risk of bias, with many studies lacking control subjects and prevalence varying between studies (Michelen et al., 2020). Differences in populations and methods make it difficult to generalize the findings across settings, particularly given the paucity of evidence from low- and middle-income countries, from primary care, and in children. Below, we describe results from some of the key studies published to date, with summaries presented in Tables 1 and 2.

A significant number of nonhospitalized patients experience persistent symptoms but estimates vary greatly among studies

The largest set of data with the right denominator to calculate prevalence for the **United Kingdom** comes from the Office for National Statistics (ONS) Coronavirus (COVID-19) Infection Survey, UK (Office for National Statistics, 2021), which has performed weekly tests for COVID-19 since April in a nationally representative sample. Simultaneously, the investigators asked a subset of 9 063 respondents who ever tested positive for COVID-19 (including those without symptoms) whether they had experienced persistent fatigue, cough, headache, loss of taste, loss of smell, myalgia, sore throat, fever, shortness of

breath, nausea/vomiting, diarrhoea or abdominal pain within 5 weeks of infection. Up until 14 December 2020, approximately 1 in 5 exhibited symptoms 5 weeks after the initial infection. Among those aged 25-69 this was 1 in 4, with similar proportions in men and women. Of these, the most common symptoms were fatigue, cough and headache, each of which were present in approximately 10% of those with symptoms at 5 weeks; 1 in 7 children aged 2-10 also reported symptoms at 5 weeks, demonstrating that this condition is not limited to adults alone. However, the ONS symptom list was limited and may have not captured those who experienced symptoms that were not listed in the questionnaire. In addition, these surveys lack control groups as some of these symptoms may occur in those who have not had COVID-19. Further data is needed to establish the clinical relevance of such findings in children.

The second largest study in the **United Kingdom** is led by researchers at King's College London who developed the COVID Symptom Study app, together with health data science company ZOE Global, which is now used by several million people (Sudre et al., 2020). They identified 4 182 people who had laboratory-confirmed COVID and continued to log symptoms comparing them to matched negative testing controls. Of cases, 1 in 8 had symptoms lasting over 28 days compared with just 1 in 50 of controls, with 1 in 20 cases continuing to experience symptoms for over 8 weeks and 1 in 50 for over 12 weeks. Those with symptoms lasting

Table 1 Summary of selected studies o	on the prevalence of Long	COVID in nonhospitalized patients
---------------------------------------	---------------------------	-----------------------------------

COUNTRY	STUDY	SAMPLE	NUMBER OF CASES INCLUDED	RESULTS
	Office for National Statistics (2020)	Population representative	8 193	 21% had symptoms 5 weeks after infection 10% had symptoms 12 weeks after infection
United Kingdom	Sudre et al. (2020)	COVID Symptom App users (out of which 14% were hospitalized)	4 182	 13% of cases had symptoms lasting 28 days after symptom onset 5% of cases had symptoms for over 8 weeks and 2% for over 12 weeks after symptom onset
	Townsend et al. (2020)	Hospital outpatients (out of which 56% were hospitalized)	127	• 52% reported persistent fatigue at 10 weeks after symptom onset
USA	Tenforde et al. (2020)	Hospital outpatients (out of which 7% were hospitalized)	292	• 35% had symptoms after a median of 16 days after testing positively for SARS-CoV-2 infection
Switzerland	Nehme et al. (2020)	Hospital outpatients	669	 About 33% of cases had symptoms 30–45 days after diagnosis
The Netherlands and Belgium	Goërtz et al. (2020) ⁶	Facebook group for coronavirus patients with persistent complaints (out of which 5% were hospitalized)	2 113	• Over 99% infected individuals did not fully recover within 12 weeks after symptom onset

^aSome of the studies reported in this table included hospitalized patients ^bIncluded suspected cases

Source: Authors

longer than 4 weeks were also more susceptible to symptom relapses (16%) than those whose symptoms resolved within 10 days (8.4%). The researchers did point out that those using their app were not necessarily representative of everyone getting COVID-19. They had more women, fewer people over 70, and fewer people from ethnic minorities than in the general population but they were able to adjust for this. However, they could not take account of the possibility that those who did experience persisting symptoms would be more or less likely to continue to report symptoms than those whose symptoms resolved.

Another study of 127 people in the **United Kingdom** (over half of whom were admitted to hospital), used a validated fatigue questionnaire and found that 52% reported persistent fatigue at 10 weeks after initial COVID symptoms (Townsend et al., 2020). There was no association with COVID-19 severity or laboratory markers of inflammation or cell turnover but females and those with a pre-existing diagnosis of depression/anxiety had higher prevalence.

In **Switzerland**, around a third of ambulatory patients had persistent symptoms of COVID-19 between 30 and 45 days after diagnosis with COVID-19, with cough and loss of taste

or smell being most common early in the disease course, and fatigue, dyspnoea and loss of taste or smell being more persistent (Nehme et al., 2020).

Many people admitted to hospital for COVID-19 experience persisting symptoms for at least 60 days after discharge

Although more patients are surviving severe COVID-19, there are limited data on outcomes after initial hospitalization. Initial studies were limited to 60 day follow-ups, but there is now data demonstrating that symptoms can last for 12 weeks and longer, and these are also reported below.

A study of hospitalized patients from the **United Kingdom** found that 68% reported persistent fatigue, 57% sleep disturbance and 32% breathlessness after an average of 60 days after discharge (Cruz et al., 2020). However, it was not possible to compare these findings with the individuals' conditions before getting COVID. Another **French** study reported persisting fatigue in 40% of hospitalized patients, also 60 days after symptom onset, with breathlessness in 30% (Carvalho-Schneider et al., 2020). Similarly,

Table 2 Summary of selected studies on the prevalence of Long COVID in hospitalized patients postdischarge

COUNTRY	STUDY	NUMBER OF CASES INCLUDED	RESULTS
Canada	Wong et al. (2020)	78	 51% had persistently reduced quality of life and 50% had shortness of breath at 12 weeks after symptom onset
France	Carvalho-Schneider et al. (2020)	130	• 40% reported persistent fatigue and 30% breathlessness at 60 days after symptom onset
Italy	Carfì, Bernabei & Landi (2020)	143	• 87% had symptoms, 55% had three or more symptoms at 60 days after discharge
United Kingdom	Cruz et al. (2020)	119	• 68% reported persistent fatigue, 57% sleep disturbance and 32% breathlessness at 60 days after discharge
	Arnold et al. (2020)	110	• 74% had persistent symptoms, typically breathlessness and fatigue and 10% had persistent anomalies on chest X-ray or respiratory function testing at 12 weeks after discharge
USA	Donnelly et al. (2020)	2 179	 19.9% were readmitted, 9.1% died and 27% were readmitted or died within 60 days after discharge
China	Huang et al. (2021)	1 733	 76% reported persistent symptoms, and 50% had residual anomalies on chest imaging 6 months after discharge

Source: Authors

143 hospitalized Italian patients, examined at a mean of 60 days since the onset of symptoms, found that only 13% were symptom-free 55% had three or more symptoms (Carfi, Bernabei & Landi, 2020).

The largest study to look at clinical outcomes among hospital patients is from the **United Kingdom** and compared clinical outcomes in 47 780 patients discharged alive with COVID-19 compared with matched controls. After a mean follow-up of 140 days, readmission rates in COVID-19 patients were 3 times higher than controls, as were new onset cardiovascular (3 times), respiratory (27 times) and diabetic (1.5 times) events, with the greatest risks observed among those over the age of 70 and in ethnic minorities (Ayoubkhani et al., 2021). Readmission rates, reasons for readmission and mortality rates were measured after hospital discharge among patients with COVID-19 in the nationwide Veterans Affairs (VA) in the USA. Within 60 days of discharge, 354 patients (19.9%) who survived COVID-19 hospitalization were readmitted, 162 (9.1%) died, and 479 (27.0%) were readmitted or died. Of those readmitted, the most common readmission diagnoses were COVID-19 (30.2%), sepsis (8.5%), pneumonia (3.1%), and heart failure (3.1%). During readmission, 22.6% were treated in intensive care, 7.1% were mechanically ventilated, and 7.9% received vasopressors (Donnelly et al., 2020).

Symptoms may last longer than 12 weeks

While a high proportion of COVID-19 patients have persistent symptoms in the short term, the ONS Infection survey and the COVID Symptom App study show that many can suffer with symptoms for as long as 12 weeks. Earlier data from the ONS survey found that among 8 193 COVID-19 cases, 1 in 10 cases were still symptomatic 12 weeks post-infection, with symptoms lasting a median of 40 days (Office for National Statistics, 2020). In addition, data collected from 2 113 (16%) members of two Facebook groups for coronavirus in the **Netherlands and Belgium** found that fewer than 1% of infected individuals fully recover within 12 weeks (Goërtz et al., 2020).

Among those who were hospitalized, an even greater proportion report symptoms at 12 weeks. In one British study of 110 patients, 74% reported persisting symptoms, typically breathlessness and fatigue, and about 10% had persisting anomalies on chest X-ray or respiratory function (Arnold et al., 2020). Similarly, a Canadian study of previously hospitalized patients, found that 51% had persistently reduced guality of life and 50% had shortness of breath 12 weeks after discharge (Wong et al., 2020). Symptoms can persist for up to 6 months, as seen in one study of 1 733 hospitalized patients in China in which many reported fatigue or muscle weakness (63%), sleep difficulties (26%) and anxiety or depression (23%). Patients who were more unwell during admission were more likely to have ongoing pulmonary complications observed in both their pulmonary diffusion capacities and abnormal chest imaging (Huang et al., 2021).

The persistence of symptoms in some patients has implications for health services and according to a survey of GPs in the **United Kingdom** in 2020, 67% were looking after patients with COVID-19 symptoms lasting longer than 12 weeks but only 23% had access to a Long COVID clinic that they could refer into (Royal College of General Practitioners, 2020).

For now the natural history on Long COVID is unknown but, on the basis of what is known about the pathology, it seems likely that it will vary considerably, with some effects, for example a consequence of thrombosis, causing organ damage permanent, others due to persisting viral infection uncertain, and some other effects likely to resolve in weeks or months.

Young children seem to be less susceptible to disease with COVID-19 but the number of children with Long COVID is uncertain

Young children (under 11 years of age) are much less likely to develop severe illness, perhaps due to their lower levels of expression of the receptors to which the virus binds. Yet there have been reports from Europe and North America of children and adolescents with COVID-19 requiring acute admission to intensive care units with a multisystem inflammatory condition with some features of Kawasaki disease and toxic shock syndrome (Moreira, 2020). These children present with acute illness accompanied by a hyperinflammatory syndrome, leading to multiorgan failure and shock. This seems to be distinct from Long COVID, although some of the mechanisms involved may be similar.

Both the ONS survey described in section 2.3 and other informal surveys among patient groups suggest that children can experience persisting fatigue, nonspecific gastrointestinal problems, sore throats, headaches, and muscle pain and weakness. Other symptoms included fevers, nausea, mood changes, rashes, dizziness, breathing difficulties and cognitive blunting.

One Swedish case series describes five children, with a median age of 12 years, who had symptoms persisting for 6–8 months (Ludvigsson, 2020). One large **United Kingdom** study of 651 children admitted to hospital with confirmed COVID-19 found that the median age was 4.6 years, 56% were male and 43% were non-White (Swann et al., 2020). Overall, 11% of children had a systematic mucocutaneous-enteric cluster of symptoms. However, data on long-term outcomes in these populations have not been reported.

Evidence is, however, fragmentary and there is an urgent need for collection of standardized data on epidemiology, including clinical presentations, severity, and outcomes (Simpson & Lokugamage, 2020). WHO has developed a preliminary case definition and case report form for multisystem inflammatory disorder in children and adolescents (WHO, 2020b).

2.4. What are the symptoms of Long COVID?

Evidence on the clinical sequalae of Long COVID is emerging and indicates a number of particular features, reflecting the increased risk of damage to multiple body systems tissues. There is still little data on the social impacts of Long COVID but available evidence points to reduced ability or even inability to work, which has implications for labour force productivity and the cost of long-term sickness absence. Data on Long COVID in health care workers and the associated social impact are so far lacking.

People suffering from Long COVID experience symptoms affecting many body systems including fatigue, breathing difficulty and fever

The investigators of the **United Kingdom**-based COVID Symptom Study identified two clusters of symptoms. People with the first cluster of symptoms suffered only fatigue, headache and upper respiratory symptoms, such as shortness of breath, sore throat, persistent cough and loss of smell (Sudre et al., 2020). The second group also had symptoms affecting other body systems, including ongoing fever and gut problems. In general, fatigue was present all the time while headaches came and went. They also found that cardiac symptoms (palpitations and tachycardia) and neurological symptoms (concentration issues, tinnitus and peripheral neuropathy) were more common among those whose symptoms persisted beyond 28 days compared with those whose symptoms resolved within 10 days.

Another study, which is a rare example of research initiated, led and reported by people affected with the condition being described, recruited 3 762 patients from 56 countries via support groups and social media although as the authors note, this approach to recruitment makes it difficult to know how representative the results are (Davis et al., 2020). The mean number of body systems affected was 9, with the most common symptoms being fatigue (77%), postexertional malaise (72%) and cognitive dysfunction and brain fog (85%). Of those who recovered within 90 days, symptoms peaked at Week 2. Gastrointestinal and skin symptoms were especially common. In this study, 86% experienced relapses following exercise, physical or mental activity. Investigators reported a third cluster of late-onset symptoms, including cardiac changes such as palpitations, neurological (including tinnitus) and skin changes.

Other studies have reported patients describing experiencing insomnia, dizziness, anxiety and depression, chills and sweats, persistent fever, body aches, and skin rashes (Patient led research for COVID-19, 2020). Patients report that their symptoms often fluctuate in intensity and frequency, with approximately 16% in the ZOE study in the **United Kingdom** reporting relapse, with new symptoms appearing at different stages of their illness.

Long COVID can also affect specific body systems including heart, lungs and brain

Turning to individual body systems, myocardial injury has attracted much attention (Puntmann et al., 2020) in part because it has been found in previously fit athletes, a group whose cardiovascular system had been examined prior to being infected with COVID-19, thus enabling comparison pre- and post-illness (Rajpal et al., 2020). Most recently, provisional data has emerged from the COVERSCAN study in the **United Kingdom** (Dennis et al., 2020), which performed serial MRI scans on a self-selected sample of 201 middle-aged, generally healthy individuals with COVID-19, of whom 70% were female, 32% were health care workers and only 18% were initially hospitalized. The investigators found evidence of mild organ impairment in heart (32%), lungs (33%), kidneys (12%), liver (10%), pancreas (17%) and spleen (6%) and reported that almost 70% of individuals have impairments in one or more organs 4 months after initial symptoms of SARS-CoV-2 infection, which was higher than age-matched healthy controls. Of note, 23% had mild systolic dysfunction and 11% had evidence of myocarditis (Dennis et al., 2020), which was associated with more severe disease.

The Hradec Králové study on Long COVID (Czech Pneumological and Phthiseological Society, ČLS JEP, 2020) in **Czechia**, examined 79 adults 3 months after their initial infection with COVID-19. The mean age was 45 and 82% were never admitted to hospital. Almost half (49%) reported having respiratory symptoms, which were most often difficulty breathing (21%), then cough (10%), fatigue (17%) or chest pain (9%). In addition, 28% of them still had a poor sense of smell after 3 months. In this study, 37% of participants had an impaired lung function (specifically, a reduced oxygen carrying capacity), while 20% had signs of residual lung damage on imaging of the chest.

There have also been case reports of new onset Type 1 diabetes, an association that is plausible as the islet cells in the pancreas have ACE-2 receptors (Rubino et al., 2020). This is currently being studied in the COVIDIAB international study (King's College London & Monash University, 2020). However, those with pre-existing cardiometabolic diseases are also at greater risk of severe COVID-19 disease and mortality (Singh et al., 2020). A recent systematic review and meta-analysis reported the proportion of new onset diabetes of 14.4% (Sathish et al., 2020) and it will be essential to ensure long-term follow-up of these patients to track the evolution of this manifestation of Long COVID.

Neurological complications of acute COVID-19 have also been documented, with the most common neurological symptoms being headaches and altered smell and taste (Padda et al., 2020). In addition, the international patientled study of Long COVID symptoms described above recorded self-reported brain fog or cognitive dysfunction in 85% of respondents (Davis et al., 2020). The most common symptoms were poor concentration (75%) and difficulty thinking (65%). Cognitive dysfunction increased over the first 3 months and was present in all age groups. Approximately 73% experienced memory impairments. A study of 509 hospitalized patients in Chicago in the USA, found that the most frequent neurologic manifestations were myalgias (44.8%), headaches (37.7%), encephalopathy (31.8%), dizziness (29.7%), altered smell (11.4%) and taste (15.9%) (Liotta et al., 2020) with between 0.2 and 1.4% of hospitalized patients suffering from strokes, movement disorders, motor and sensory deficits, ataxia and seizures. Neurological complications were more common in those who were younger and had severe COVID-19, while those with encephalopathy were slower to regain their functional capacity and had a higher mortality than those

without. There is also some evidence of SARS-COV-2 invasion leading to persisting inflammation of the olfactory mucosa in those with continuing loss of the sense of smell (De Melo et al., 2020). However, methodological differences between different studies make it difficult to determine how common these manifestations actually are among the overall population who have COVID-19 (Pezzini & Padovani, 2020).

As with diabetes, the association with psychiatric disorders is likely bidirectional. Of GPs surveyed by the Royal College of General Practitioners in the United Kingdom, 76% described patients reporting sleep disorders or mood changes (Royal College of General Practitioners, 2020). Electronic health record data from 69 million individuals in the United Kingdom also showed that a diagnosis of COVID-19 was associated with an increased incidence of a first psychiatric diagnosis between 14 and 90 days after diagnosis (Taquet et al., 2020), with the greatest risk observed for anxiety disorders, insomnia and dementia. Similarly, those with a psychiatric diagnosis in the previous year were reported to be at a 65% greater risk of being diagnosed with COVID-19, suggesting that mental ill health could be both a risk factor and a complication of COVID-19, although this did not account for crucial differences in socioeconomic status.

Long COVID can be severely disabling with implications for people's ability to work and social lives

There is still little data on the social impacts of Long COVID, but evidence from 150 patients with non-critical COVID-19 who were discharged from hospital in France between March and June 2020 showed that 11% were still on sick leave at Day 60 (Yelin et al., 2020). Half of the cohort were health care workers, posing a reminder of the risks that Long COVID presents to health systems. Similarly, the international patient-led survey described above found that 45% of Long COVID sufferers had reduced their workload and 22% were not working due to their illness (Davis et al., 2020). Similar findings were observed in Canada when SARS took hold and researchers at the University of Toronto described 22 people with SARS who were unable to work for up to 36 months after initial infection (Moldofsky & Patcai, 2011). Previous experience with chronic fatigue syndrome (CFS) has also taught us a great deal about the potential costs of long-term sickness absence (Reynolds et al., 2004). In one study from the USA, household productivity declined by 37% and labour force productivity by 54% among people with CFS. The annual lost productivity in the USA was US\$ 9.1 billion, representing about US\$ 20 000 per person with CFS. This accounts for around half of the household and labour force productivity of the average person with this syndrome. Data on Long COVID in health care workers and social impacts are lacking although a group of health care workers have documented their experiences of COVID-19 and persistence of symptoms and have made recommendations about potential methods of follow-up and Long COVID surveillance (Alwan et al., 2020). A UK Doctors Long COVID group has also been set

up and has reported on symptoms they experience, including myocarditis or pericarditis, microvascular angina, cardiac arrhythmias including atrial flutter and atrial fibrillation, dysautonomia including postural orthostatic tachycardia syndrome, mast cell activation syndrome, interstitial lung disease, thromboembolic disease (pulmonary emboli or cerebral venous thrombosis), myelopathy, neuropathy and neurocognitive disorders, renal impairment, new onset diabetes and thyroiditis, hepatitis and abnormal liver enzymes, new onset allergies and anaphylaxis and dysphonia, all of which impact on their ability to continue working (Gorna et al., 2020).

2.5. Diagnosis and treatment

There is no simple symptom or test for diagnosing Long COVID

Long COVID is a clinical diagnosis, based on a history of COVID-19 and a failure to fully recover, with development of some of the symptoms listed in section 2.4. Although a positive swab or antibody test for COVID-19 is helpful, it is not a prerequisite for diagnosis. This is because availability of testing was very limited for to most people in Europe in the early stages of the pandemic. Moreover, PCR testing is not 100% sensitive and both it and antibody testing may miss those with residual symptoms from older infections, as antibodies often decline substantially by about 4 months (Gudbjartsson et al., 2020).

Long COVID is different from the well-recognized postintensive care syndrome, which is marked by chronically impaired pulmonary function, neuromuscular weakness, long-term psychological impacts and reduced quality of life (Ahmed et al., 2020; Society of Critical Care Medicine, 2020), a pattern that was also observed in SARS survivors (Ahmed et al., 2020). The syndrome is common among people with severe acute infections who have spent long periods on ventilators (Desai, Law & Needham, 2011) and may coexist with Long COVID in some cases.

While the symptoms of Long COVID do bear some semblance to the postviral symptoms observed in those with CFS (Institute of Medicine, 2015), Long COVID seems to manifest as a broader spectrum of symptoms and may potentially be more common, although further study is required to confirm this (Perego et al., 2020).

Emerging evidence points to Long COVID having a number of particular features, at least some of which seem to be related to the increased risk of damage to the certain tissues, in particular cardiac, vascular, lung, kidney involvement with evidence implicating both infection by the virus and the resulting inflammatory response (see Box 1).

In the United Kingdom, the National Institute for Clinical Excellence (NICE) have produced guidelines, in which they define some criteria for classifying these ongoing symptoms, which are presented in Box 2.

Box 2: NICE guidelines on diagnosing Long COVID

Guidelines developed by NICE in England (National Institute for Health and Clinical Excellence, 2020) advise the following criteria for diagnosing persistent symptoms of COVID-19:

- ongoing symptomatic COVID-19 if people present with symptoms 4 to 12 weeks after the start of acute COVID-19 that are not explained by an alternative diagnosis; or
- **post-COVID-19 syndrome** if the person's symptoms have not resolved 12 weeks after the start of acute COVID-19.

Unfortunately, the guidelines do not specify which symptoms should be included in this definition, which could introduce unnecessary variability, but they do recommend that investigation and management should be based in primary care with referral of more complex cases to specialist post-COVID clinics.

In response to the NICE guidance, doctors and patient groups in the **United Kingdom** have called for clearer case definitions and comprehensive descriptions of the variety of known symptoms of Long COVID to ensure health care workers are able to identify cases early (Gorna et al., 2020). They argue that given that the underlying immunological and organ specific effects of SARS-COV-2 are now better understood, this should not be considered a diagnosis of exclusion (Sivan & Taylor, 2020). This would enable prompt assessment and earlier diagnosis, as well as better treatment and rehabilitation.

A multidisciplinary approach to assessment and management of Long COVID is essential

Given the complex nature of the condition, with implications for many aspects of the lives of those affected, NICE guidelines advise that assessment and management should be tailored to the individual's problems, after excluding any coexisting illness that may be giving rise to the symptoms reported. For example, this would include a chest X-ray in those with respiratory symptoms, checking for a postural drop in blood pressure in those with dizziness, and urgent referral to the appropriate specialist for those with, for example, chest pain, palpitations or mental illness. Those who need further care should be referred to multidisciplinary rehabilitation teams (using screening questionnaires such as C19-YRS (Yorkshire Rehabilitation Screen)) (Sivan, Halpin & Gee, 2020) or for specialist input in more complex cases of specific organ dysfunction. The implications for health services of these guidelines are discussed further below (see section 3.2).

3. Policy and other responses in Europe

In this section we list a number of responses to Long COVID that have been adopted in different European countries. To that end, we contacted participants in the European Observatory COVID-19 Health Systems Response Monitor and undertook a rapid Internet search. The following examples should not be considered exhaustive, which would be impossible in this rapidly moving field, but do serve to illustrate some of the emerging ideas being considered. It

should also be noted that while the text was checked with national correspondents at the time of publication, it is to be expected that further developments will continue to take place.

3.1. Organized national surveillance systems

Surveillance is a critical part of monitoring Long COVID but is not widespread in Europe

Surveillance of Long COVID is particularly important but is not happening routinely in European countries. A first step in implementing a surveillance system is to define and classify it (Alwan & Johnson, 2020). To support this process, WHO has now introduced post-COVID emergency use ICD codes, which include "Personal history of COVID-19", "Post COVID-19 condition", and "Multisystem inflammatory syndrome associated with COVID-19" (WHO, 2020b) (Box 3).

Box 3: ICD-10 codes for use with Long COVID

U08.9 Personal history of COVID-19, unspecified

Note: This optional code is used to record an earlier episode of COVID-19, confirmed or probable that influences the person's health status, and the person no longer suffers of COVID-19.

U09.9 Post COVID-19 condition, unspecified

Note: This optional code serves to allow the establishment of a link with COVID-19.

This code is not to be used in cases that still are presenting COVID-19.

U10.9 Multisystem inflammatory syndrome associated with COVID-19, unspecified

Temporally associated with COVID-19:

- Cytokine storm
- Kawasaki-like syndrome
- Paediatric Inflammatory Multisystem Syndrome (PIMS) With COVID-19
- Multisystem Inflammatory Syndrome in Children (MIS-C)
- Excludes: Mucocutaneous lymph node syndrome [Kawasaki] (M30.3)

Although most countries now have COVID-19 data surveillance systems that record the total number of positive diagnoses, deaths, hospital admissions and intensive care unit (ICU) admissions (Alwan, 2020b) we were unable to find any details of specific data streams for Long COVID, although some, such as those in Italy (Istituto Superiore di Sanità, 2020) and Czechia (Ministry of Health of the Czech Republic, 2020), record the proportion of patients who recover, although this usually depends on returning a negative test rather than being free of symptoms (Alwan, 2020a).

3.2. Organized health systems responses

Patients with Long COVID often report their difficulties are not taken seriously

Patients suffering with Long COVID have reported feeling stigmatized and unable to access and navigate services. In particular, they report difficulties in being taken seriously and achieving a diagnosis. The care they received was reported to be disjointed and siloed and specialist care was mostly inaccessible and variable. While some felt supported, many report feeling "fobbed off" (Ladds et al., 2020).

Despite this, health systems in several countries are now developing organized responses.

National health system responses include development of dedicated treatment guidelines and pathways, and creation of post-COVID clinics and online support tools

As mentioned above, in **England**, NICE, working with patient groups and professional associations, has published a set of guidelines (National Institute for Health and Clinical Excellence, 2020). They propose multidisciplinary assessment services, bringing together physicians with expertise in different body systems, as well as multidisciplinary rehabilitation services, with core teams that could include, but not be limited to, occupational therapy, physiotherapy, clinical psychology and psychiatry, and rehabilitation medicine. These services should be part of a model of care that includes integrated referral patterns between primary, secondary and community care.

Prior to the publication of these guidelines the English National Health Service (NHS) had set out its own care pathway for patients with ongoing symptoms persisting beyond 6 weeks (NHS, 2020a). It stipulated that an initial assessment should be undertaken in primary or secondary care to exclude serious underlying pathology, using a screening tool included in the documentation. Onward referral would then be negotiated by a single point of access to triage and refer on to one or more of three possible pathways for those whose symptoms persist beyond 12 weeks: post-COVID assessment clinics, local rehabilitation clinics or the "Your COVID Recovery" NHS digital programme (https://www.yourcovidrecovery.nhs.uk). The NHS announced the launch of 40 such post-COVID assessment clinics in England in Autumn 2020, which adopt standardized functional assessment and screening questionaries coupled with secondary care specialist referrals for investigation of potential end organ sequelae, where suspected. Physical care is also integrated with mental health care provision, including self-care and self-help groups, community therapy where required and referral to the "Your COVID Recovery" NHS digital programme to support COVID recovery (NHS, 2020b). This digital online platform includes self-management advice on tackling daily activities, returning to work and seeking help. All cases of acute COVID-19 admitted to hospital will also be reviewed remotely and face to face at 6 weeks for a review of symptoms to identify cases for onward referral. It will be important for these clinics to have standard operating procedures for assessment, documentation and long-term

follow-up data of those managed in these clinics. A number of other services have also appeared including online Long COVID support groups (Long COVID Support, 2020), and NHS resources (NHS, 2020a).

In **Germany** some larger university hospitals offer special consultation hours for patients with long-term complaints, including the Jena University Hospital or Hanover Medical School, which opened a post-COVID outpatient clinic in in July 2020 where neurologists, cardiologists, pneumologists, psychiatrists, gastroenterologists, occupational physicians, internists and ear, nose and throat doctors provided interdisciplinary care. Unfortunately, access to such clinics was not widespread at the time of writing.

In Italy, service delivery for COVID-19 varies regionally but in general is managed in primary and secondary care and facilitated through community care networks, including a combination of reconverted community hospitals, health care hotels, nursing homes and low intensity residential facilities. GPs also coordinate services that include domiciliary care, supported by remote telemonitoring and a new community nurse role. Some hospitals, such as the San Raffaele Hospital in Lombardy, have also created post-COVID wards for rehabilitation of those who experienced prolonged admissions to ICU. Other services include those by AbilityAmo (AbilityAmo, 2020), a non-profit organization, which is dedicated to providing rehabilitative interventions for post-COVID disability and fragility. These services include telemonitoring systems for clinical surveillance, specialist interventions (both in hospitals and at home), psychological support and neurological, respiratory and cognitive postadmission rehabilitation. The organization is also gathering information about patients' experiences and plans to publish a report on life after COVID-19.

In Czechia, the Hradec Králové teaching hospital set up a specialized Post-COVID Care Centre to take care of Long COVID patients with residual symptoms 3 months after their acute infection. The care centre provides a multidisciplinary team of physicians, including pneumologists, cardiologists, neurologists, psychiatrists, nephrologists, dermatologists, gastroenterologists, rheumatologists and rehabilitation physicians. The intention is for patients to be cared for by their GPs, with follow-ups by respiratory doctor until symptoms and any residual organ damage have resolved (Biba, 2021). This particular centre primarily serves the Královehradecký and Pardubický regions but others are also expected to open elsewhere. The Czech Pneumological and Phthiseological Society of the J.E. Purkyně Czech Medical Association (ČLS JEP) also published interim guidance in early January 2021, including a definition, diagnosis and classification of Long COVID and post-COVID syndrome (Czech Pneumological and Phthiseological Society, ČLS JEP, 2021). Patients with ongoing symptoms such as breathlessness have been advised by the Czech Pneumological and Phthiseological Society of the ČLS JEP to seek help from a respiratory doctor, but no specific services seem to have been designed.

Similarly, in **Spain**, hospitals and primary care centres provide COVID-19 patients with guidance on respiratory and physical rehabilitation to face the consequences of COVID-

19 and the Spanish Society of General Practitioners has published guidelines about treating this type of patients in primary care settings (Sociedad Española de Médicos Generales y de Familia, 2020a).

In Belgium, France and Portugal, specific services are mainly developed for COVID patients who were hospitalized. In Belgium, services are focused on patients who were admitted to intensive care units, with programmes dedicated to the management of the postintensive care syndrome (PICS). These services can take place at home, in a specialized or a geriatric hospital and include a multidisciplinary team of doctors, nurses, physiotherapists, occupational therapists, sports therapists, psychologists, speech therapists and social workers. As well as neuromuscular rehabilitation, there are also functional rehabilitation centres for locomotor and neurological disorders, with cognitive rehabilitation interventions provided where necessary. The COVID-19 pandemic has also accelerated the expansion of pre-existing intensive post-care pathways. At the time of writing, at least three Belgian hospitals were developing similar care pathways, while others were considering it. Some hospitals have also dedicated specific care units to rehabilitate these patients, given the limitations of infection control measures for COVID positive patients in hospital.

Responses to Long COVID are also developed by local service providers and professional bodies

There are also many examples of local initiatives. Thus, the Fondazione Policlinico Agostino Gemelli in Italy, a tertiary care centre in Rome, has developed a multidisciplinary health care service called the Post-COVID-19 Day-Hospital, which includes an infectious diseases physician and specialists for pneumological assessments, ophthalmologic assessments, otolaryngologic assessments, neurologic assessments, psychiatric assessments, cardiovascular assessments, nutritional assessments, gut assessment and rheumatologic assessments (Landi et al., 2020). Some hospitals in the United Kingdom have also offered information packs with recommendations (Homerton University Hosptal, 2020) to improve breathlessness, diet, emotional well-being and conserving energy such as breathing techniques, physical position and relaxation. There are also initiatives by professional bodies. Thus, occupational therapists in the United Kingdom have also released guidance on how to conserve energy (Royal College of Occupational Therapists, 2020), while the Society for Critical Care Medicine has released guidance on PICS and how to manage it.

Services that have been developed in many European countries so far are not universally accessible or multidisciplinary and many do not provide access to disability benefits

We were unable to find many examples of universally accessible multidisciplinary services for Long COVID that were available to communities in European countries outside the United Kingdom. Similarly, few countries had specific disability benefits for patients with Long COVID or any mechanisms designed to protect employment in these individuals, although there is some guidance from the Faculty of Occupational Medicine in the United Kingdom to help health care workers support Long COVID patients back to work.

3.3. Patient involvement

Patients have called for recognition, rehabilitation and research

The patient voice has been critical in shaping awareness about Long COVID (Callard & Perego, 2021), although to a greater extent in some countries than in others. A number of very popular patient advocacy groups have been launched on social media across Europe and in other countries, including the Body Politic group (Body Politic, 2020a), which also conducted some of the first patient-led research to document and raise awareness of the symptoms among sufferers, many of whom describe feeling unheard by doctors (Jensen, 2020).

The United Kingdom arm of Body Politic has set up LongCovidSOS (Body Politic, 2020b), an advocacy group that has appealed to governments to recognize and raise awareness of the needs of Long COVID sufferers among the public, and employers. They called for "Recognition, Research and Rehabilitation"; a call supported by patient associations in Belgium and that has been publicly acknowledged by the Director-General of WHO.

- Recognition involves taking into consideration all aspects of the problem (medical, psychological, social, etc.) and raising the awareness among physicians and other health professionals (physiotherapists, nurses, etc.) about Long COVID.
- Rehabilitation involves the provision of nationally coordinated multidisciplinary programmes for Long COVID patients to assess, test, diagnose and care, including those who were not hospitalized during the acute phase.
- Research calls for coordinated and co-created multidisciplinary studies to understand the clinical sequelae and develop treatment pathways for Long COVID.

Patient groups have also asked authorities to set up working groups to address the needs of Long COVID sufferers; and focus on the economic implications, including the provision of financial and employment support.

In **Spain**, several Long COVID online communities have been formed in collaboration with the Spanish Society of General Practitioners (SEMG) to raise awareness of their condition and improve health care services (Sociedad Española de Médicos Generales y de Familia, 2020b). In France, patients initially organized themselves in social media groups but as of 13 October 2020 they created an association (#AprèsJ20-Association Covid long France) to try to coordinate their advocacy actions (Association COVID Long France, 2020). Several Facebook groups have been created for doctors suffering from Long COVID, many of whom became infected in the first wave of the pandemic when personal protective equipment (PPE) was in short supply. Some have appealed to the medical profession to recognize their needs, calling for collaboration between politicians, health care services, public health professionals, scientists and society to improve our understanding of the epidemiology, pathophysiology and management of Long COVID, highlighting the need to measure and quantify Long COVID (Alwan et al., 2020). They have also called for timely access to patient-centred services that can investigate and treat pathology and focus on supporting functional recovery, whether or not an individual has been tested for COVID-19.

3.4. Self-care and self-help groups

Self-help groups have been created in many countries and have helped to articulate the patient voice and provide support

In Germany, the earliest Long COVID self-help groups started in March 2020, in hardest-hit areas in North Rhine-Westphalia (e.g. Heinsberg) and Bavaria. Participants use the forums to exchange experiences of the disease, to support each other and to gain new perspectives. Italy has also seen a number of patient initiatives, including a social network group called "We who have beaten COVID" (Noi che il Covid l'abbiamo sconfitto) (Noi che il Covid lo abbiamo sconfitto, 2020) with 12 630 subscribers at the beginning of January 2021. Belgium has also seen many patients, including health workers, join social media groups in both French and Flemish languages, with some hospitals also setting up their own discussion groups, many of whom are now contributing to research studies into clinical features and management of Long COVID. A group focusing on Long COVID in children (https://www.longcovidkids.org/) has been created in the United Kingdom.

Initiatives have also been created in some countries to promote self-care among those with Long COVID, to complement existing services

In addition to Facebook groups in the United Kingdom (some of which have tens of thousands of members), Asthma UK and the British Lung Foundation have come together with the support of others including the British Thoracic Society and the Primary Care Respiratory Society to develop the online Post-COVID HUB (https://www.postcovid.org.uk) for people left with breathing difficulties following a COVID-19 infection to access advice and support to manage symptoms (Asthma UK and British Lung Foundation, 2020), while others have united to create selfhelp pacing groups, supporting each other to monitor and very slowly increase their daily activity within their limits (Hammond et al., 2020). In **Portugal**, the Ministry of Health and the Directorate General of Health have developed a series of online materials targeting mental health of health professionals (Direção Geral da Saúde, 2020a) including a guide for self-care and mental health (Direcão Geral da Saúde, 2020b).

3.5. Ongoing research

Studies of Long COVID have been emerging in Europe, offering opportunity to improve our understanding of this new condition

The COVID-19 Research Project Tracker (UKCDR, 2020) is a live database of funded research projects across the world related to the current COVID-19 pandemic that has been set up by the United Kingdom Collaborative on Development Research (UKCDR) and the Global Research Collaboration For Infectious Disease Preparedness (GLOPID-R) to map research initiatives on to the WHO's Coordinated Global Research Roadmap for COVID-19.

In the **United Kingdom**, a number of studies are already up and running in relation to Long COVID, and the United Kingdom Research and Innovation (UKRI) and the National Institute for Health Research (NIHR) have also launched research calls for up to £ 20 million to fund several others in 2021 (UKRI, 2020). Existing research is ongoing in the United Kingdom, including the Post-Hospitalisation COVID-19 study (PHOSP-COVID, 2020), which aims to follow-up inpatients with COVID-19 after discharge to document their symptoms, blood markers, imaging and patient reported outcomes. The research team from Leicester have recruited 1 500 people so far and plan to recruit a further 8 500, with an additional aim planned in 2021 to include people with COVID-19 who were not admitted to hospital during the acute phase. A number of subgroups of PHOSP-COVID study have also been set up to conduct research in specific groups including cardiovascular, metabolic, multimorbidity and ethnic subgroups. In addition, working groups from the multidisciplinary Long COVID community clinics will investigate the long-term consequences of COVID-19 and their mechanisms, the impact of acute interventions, and identify future research and intervention needs. These groups will be exploring answers to research questions emerging from the social sciences and the health economic impacts of Long COVID. The PHOSP-COVID study is also linked to other immunology and neuropsychology studies in the United Kingdom and to the International Severe Acute Respiratory and emerging Infection Consortium (ISARIC) Global COVID-19 Long-Term Follow-Up study (University of Oxford, 2020), which has developed standardized data collection tools to document the clinical sequelae and risk factors for longer term consequences of COVID-19 for up to 3-5 years and to facilitate international comparisons in adults and children. UK_REACH will be the largest study of prevalence of COVID and Long COVID, complications and wider social impacts on health care workers in the United Kingdom so far. The study will include routinely collected data and will conduct long-term follow-up of 30 000 health care workers (NIHR Leicester Biomedical Research Centre, 2020).

Similar studies are emerging across Europe; for example, in Lombardy, Italy, where researchers are also studying individuals discharged from hospital with COVID-19, while the Spanish Society of General Practitioners has also launched an online anonymous survey to explore ongoing symptomatology in COVID-19 patients. As part of the Network of University Medicine in Germany, the National Pandemic Cohort Network at University Hospitals (NAPKON) investigates long-term organ damage and subsequent morbidities after infection with SARS-CoV-2 across all disease severities through population-based cohort studies at several sites (ClinicalTrials.gov, 2020a), while others in Munich, Germany, are also examining lung function to help to establish treatment protocols and to target follow-up care.

Another study at the University of Tübingen investigates the long-term consequences of individuals with mild courses of COVID-19 who have never been admitted to hospital due their illness. The researchers closely collaborate with GPs and public health offices. Finally, the Hradec Králové study on Long COVID from Czechia (described in section 2.4) also plans to repeat all physical examinations twice over the course of the year to establish whether lung function resolves in sufferers of Long COVID and will assess antibody levels and cell-mediated immunity, which is also being examined in the COVIMMUNE study in France (ClinicalTrials.gov, 2020b). There is a hope that the use of standardized methods will facilitate the production of a meta-analysis as findings from these studies start to emerge. Similarly, in Belgium, researchers at the Health Care Knowledge Centre (KCE) are also are working to establish the clinical segualae and risk factors. They are also investigating the experiences and unmet needs of patients.

4. Implications for policy

Although Long COVID is not yet fully understood, health policy-makers should be preparing to address it

Although there are still many unanswered questions about the natural history of Long COVID, in particular relating to its likely duration in those affected, it is probable that there will be considerable numbers of people with ongoing symptoms for several years at least. Health policy-makers should be preparing responses to those with this condition.

First step towards dealing with Long COVID should include implementing effective patient registers or other surveillance systems

A first step will be to ensure that there are effective systems of surveillance in place, ideally creating patient registers which are not entirely dependent on diagnosis by lab confirmation and including these figures within reported outcome measures for COVID-19. Identification, monitoring and recording of people with Long COVID is imperative to facilitate long-term follow-up and prevent an epidemic of chronic disease with detrimental impacts on health systems and economies and can also act as a critical basis for research. These registries should collect sufficient data to characterize factors that influence the risk of ongoing problems, including initial presentation and sociodemographic data, including ethnicity and occupation. They should also monitor the impact of this condition on the lives of those affected, including conventional measures of quality of life but also the impact on psychological condition, employment and economic situation.

Care guidelines and multidisciplinary services need to be developed to ensure appropriate assessment and management of the condition

In countries where this has not yet happened, it will be important to develop contextually appropriate guidelines for health professionals caring for those affected, especially in primary care. As described in this policy brief, there is already considerable material that can be drawn upon. Patient groups have emphasized that new nomenclature must not be developed by each individual guideline, but that a proper consultative process occurs by WHO to determine terminology. This will ensure consistency in diagnosis across different regions of the world and therefore better tracking of the impact of COVID on long-term morbidity.

In parallel, it will be necessary to develop multidisciplinary and multispecialty services that can evaluate individuals and ensure that they have access to appropriate management. There is considerable scope for implementation research and, as knowledge accumulates, evaluation of different forms of treatment, ideally in clinical trials. This is an area that will benefit from international cooperation.

Effective response can only be achieved by involving Long COVID patients themselves

Patient groups have already proven themselves to be instrumental to recognizing, characterizing this condition and advocating for it to be properly and fairly managed. Again, there are many opportunities for cross-country learning and such initiatives should be welcomed, encouraged, and where possible, supported by policy-makers.

References

- AbilityAmo (2020). COVID-19 EMERGENZA. Turin: AbilityAmo (http://abilityamo.it/campaign/covid-19emergenza/, accessed 28 January 2021).
- Ahmed H et al. (2020). Long-term clinical outcomes in survivors of severe acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after hospitalisation or ICU admission: A systematic review and meta-analysis. J Rehabil Med.52(5):jrm00063.
- Altmann DM, Boyton JR (2020). Confronting the pathophysiology of long covid. *BMJ Opin.* 9 December 2020.
- (https://blogs.bmj.com/bmj/2020/12/09/confronting-thepathophysiology-of-long-covid/?fbclid=IwAR0ov1rCW2T9 DbJtXs91tFVrMc9H2kTbInNJ1QdmT90rQCXPqtYI5kjPMt4, accessed 28 January 2021).
- Alwan NA (2020a). A negative COVID-19 test does not mean recovery. *Nature*.584:170.
- Alwan NA (2020b). Surveillance is underestimating the burden of the COVID-19 pandemic. Lancet.396(10252):e24.
- Alwan N, Johnson L (2020). Long COVID: where do we start with the case definitions? *OSF Preprints*. (https://osf.io/hndtm/, accessed 28 January 2021).
- Alwan NA et al. (2020). From doctors as patients: a manifesto for tackling persisting symptoms of covid-19. BMJ.370:m3565.
- Arnold DT et al. (2020). Patient outcomes after hospitalisation with COVID-19 and implications for followup: results from a prospective UK cohort. *Thorax*. doi: 10.1136/thoraxjnl-2020-216086.
- Association COVID Long France (2020). *Qui sommes nous?* (https://www.apresj20.fr/qui-sommes-nous, accessed 28 January 2021).
- Asthma UK and British Lung Foundation (2020). Post-COVID HUB. London: Asthma UK and British Lung Foundation Partnership (https://www.post-covid.org.uk/, accessed 28 January 2021).
- Ayoubkhani D et al. (2021). Epidemiology of post-COVID syndrome following hospitalisation with coronavirus: a retrospective cohort study. *medRxiv*.2021.01.15.21249885. doi: 10.1101/2021.01.15.21249885.
- Bellini WJ et al. (2005). Subacute sclerosing panencephalitis: more cases of this fatal disease are prevented by measles immunization than was previously recognized. J Infect Dis.192(10): 1686–93.
- Biba L (2021). Nemocnice v Hradci Králové otevírá postcovidové centrum. Bude pečovat o pacienty s přetrvávajícími symptomy. Hospodářské Noviny. 19 January 2021 (https://domaci.ihned.cz/c1-66870940nemocnice-v-hradci-kralove-otevira-postcovidove-centrumbude-dlouhodobe-pecovat-o-pacienty-s-pretrvavajicimisymptomy, accessed 28 January 2021).
- Body Politic (2020a). COVID-19 support group (https://www.wearebodypolitic.com/covid19, accessed 28 January 2021).

- Body Politic (2020b). LongCovidSOS
- (https://www.longcovidsos.org/, accessed 28 January 2021).
- British Society for Immunology (2020). Long-term immunological health consequences of COVID-19. London: British Society for Immunology (https://www.immunology.org/sites/default/files/BSI_Briefin g_Note_August_2020_FINAL.pdf, accessed 28 January 2021).
- Callard F, Perego E (2021). How and why patients made Long Covid. *Soc Sci Med*.268:113426. doi: 10.1016/j.socscimed.2020.113426.
- Carfi A, Bernabei R, Landi F (2020). Persistent symptoms in patients after acute COVID-19. *JAMA*.324(6):603–5.
- Carvalho-Schneider CE et al. (2020). Follow-up of adults with noncritical COVID-19 two months after symptom onset. *Clin Microbiol Infect*.S1198-743X(20)30606-6. doi: 10.1016/j.cmi.2020.09.052.
- ClinicalTrials.gov (2020a). COVIDOM: Longterm morbidity of SARS-CoV-2 infection and COVID-19 disease – consequences for health status and quality of life (NAPKON-POP). Bethseda, MD: National Library of Medicine (https://clinicaltrials.gov/st2/chow/NCT04670584_accessed
- (https://clinicaltrials.gov/ct2/show/NCT04679584, accessed 28 January 2021).
- ClinicalTrials.gov (2020b). Study of the incidence of SARS-CoV-2 infection (COVID-19) (CovImmune 2). Bethseda, MD: National Library of Medicine (https://clinicaltrials.gov/ct2/history/NCT04429594?V_2=Vi ew, accessed 28 January 2021).
- Cruz RF et al. (2020). Chest radiography is a poor predictor of respiratory symptoms and functional impairment in survivors of severe COVID-19 pneumonia. *ERJ Open Res*.00655–02020. doi: 10.1183/23120541.00655-2020.
- Czech Pneumological and Phthiseological Society, ČLS JEP (2020). Vyléčené z koronaviru by měl sledovat plicní lékař. Nemoc totiž může zanechat stopy. Prague: Czech Pneumological and Phthiseological Society (http://www.pneumologie.cz/novinka/1670/vylecene-zkoronaviru-by-mel-sledovat-plicni-lekar-nemoc-totiz-muzezanechat-stopy/ accessed 28 January 2021).
- Czech Pneumological and Phthiseological Society, ČLS JEP (2021). Post-Covid syndrome – definition, diagnosis, classification; position document. Prague: Czech Pneumological and Phthiseological Society (http://www.pneumologie.cz/upload/1610134989.5492.d ocx, accessed 28 January 2021).
- Davis HE et al. (2020). Characterizing Long COVID in an international cohort: 7 months of symptoms and their impact. *medRxiv*.2020.12.24.20248802. doi: 10.1101/2020.12.24.20248802.
- De Melo GD et al. (2020). COVID-19-associated olfactory dysfunction reveals SARS-CoV-2 neuroinvasion and persistence in the olfactory system. *bioRxiv*:2020.2011.2018.388819. doi: 10.1101/2020.11.18.388819.
- Dennis A et al. (2020). Multi-organ impairment in low-risk individuals with long COVID. *medRxiv*.2020.2010.2014.20212555. doi: 10.1101/2020.10.14.20212555.

Desai SV, Law TJ, Needham DM (2011). Long-term complications of critical care. Crit Care Med. 39(2):371–9.

Direção Geral da Saúde (2020a). Ajudar os profissionais de saúde durante o surto de COVID-19. Lisbon: Direção Geral da Saúde (https://saudemental.covid19.min-saude.pt/ajuda-a-profissionais-de-saude/, accessed 28 January 2021).

Direção Geral da Saúde (2020b). COVID-19 – Autocuidado e bem-estar dos profissionais de saúde durante a pandemia. Lisbon: Direção Geral da Saúde (https://www.dgs.pt/documentos-e-publicacoes/guia-deautocuidado-e-bem-estar-dos-profissionais-de-saude-dura nte-a-pandemia-pdf.aspx, accessed 28 January 2021).

Donnelly JP et al. (2020). Readmission and death after initial hospital discharge among patients with COVID-19 in a large multihospital system. *JAMA*.325(3):304–6. doi: 10.1001/jama.2020.21465.

Goërtz YMJ et al. (2020). Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? *ERJ Open Res*.6(4):00542-2020. doi: 10.1183/23120541.00542-202.

- Gorna R et al. (2020). Long COVID guidelines need to reflect lived experience. *Lancet*.S0140-6736(20)32705-7. doi: 10.1016/S0140-6736(20)32705-7.
- Greenhalgh T et al. (2020). Management of post-acute covid-19 in primary care. *BMJ*.370:m3026.

Gudbjartsson DF et al. (2020). Humoral immune response to SARS-CoV-2 in Iceland. N Engl J Med.383(18):1724–34.

Gupta A et al. (2020). Extrapulmonary manifestations of COVID-19. *Nature Med*.26(7):1017–32.

Hammond I et al. (2020). Long covid and self-help pacing groups—getting by with a little help from our friends. *BMJ Opin.* 29 September 2020 (https://blogs.bmj.com/bmj/2020/09/29/long-covid-andself-help-pacing-groups-getting-by-with-a-little-help-fromour-friends/, accessed 28 January 2021).

Haute Autorité de Santé (2021). Covid-19: diagnostiquer et prendre en charge les adultes présentant des symptômes prolongés. (https://www.has-sante.fr/jcms/p_3237458/fr/covid-19-diagnostiquer-et-prendre-en-charge-les-adultes-presentant-des-symptomes-prolonges, accessed 17 February 2021).

Homerton University Hosptal (2020). Post COVID-19 patient information pack. London: Homerton University Hospital NHS Trust (https://www.hackneycitizen.co.uk/wpcontent/uploads/Post-COVID-19-information-pack-5.pdf? fbclid=lwAR0g_r58IMe9nKjFfiBiGY0_9VZ0_h-08MiBViRGBS3yLelHxPkaSRXW6cc, accessed 28 January 2021).

Huang C et al. (2021) 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. Lancet.397(10270):220–32. doi: 10.1016/S0140-6736(20)32656-8.

Institute of Medicine (2015). Beyond myalgic encephalomyelitis/chronic fatigue syndrome: redefining an illness. Washington DC, National Academies Press.

Istituto Superiore di Sanità (2020). COVID-19 integrated surveillance data in Italy. Rome: Istituto Superiore di Sanità (https://www.epicentro.iss.it/en/coronavirus/sars-cov-2dashboard, accessed 28 January 2021). Jensen L (2020). "Just stay at home" was lonely and terrifying. *BMJ*.371:m3807.

King's College London, Monash University (2020). COVIDIAB registry (http://covidiab.e-dendrite.com/, accessed 28 January 2021).

Ladds E et al. (2020). Persistent symptoms after Covid-19: qualitative study of 114 "long Covid" patients and draft quality principles for services. *BMC Health Serv Res*.20(1):1144.

Landi F et al. (2020). Post-COVID-19 global health strategies: the need for an interdisciplinary approach. *Aging Clin Exp Res*.32(8):1613–20.

Libby P, Lüscher T (2020). COVID-19 is, in the end, an endothelial disease. *Eur Heart J*.41(32):3038–44.

Liotta EM et al. (2020). Frequent neurologic manifestations and encephalopathy-associated morbidity in Covid-19 patients. *Ann Clin Transl Neurol*.7(11):2221–30.

Long COVID Support (2020). Resources (https://www.longcovid.org/resources, accessed 28 January 2021).

Ludvigsson JF (2020). Case report and systematic review suggest that children may experience similar long-term effects to adults after clinical COVID-19. Acta Paediatr. doi: 10.1111/apa.15673.

Michelen M et al. (2020). Characterising long-term covid-19: a rapid living systematic review. *medRxiv*.2020.2012.2008.20246025. doi: 10.1101/2020.12.08.20246025.

Ministry of Health of the Czech Republic (2020). COVID 19: Přehled aktuální sítuace v ČR. Prague: Ministry of Health of the Czech Republic (https://onemocneniaktualne.mzcr.cz/covid-19, accessed 28 January 2021).

Moldofsky H, Patcai J (2011). Chronic widespread musculoskeletal pain, fatigue, depression and disordered sleep in chronic post-SARS syndrome; a case-controlled study. *BMC Neurol*.11:37.

Moreira A (2020). Kawasaki disease linked to COVID-19 in children. *Nat Rev Immunol*.20(7):407.

National Institute for Health and Clinical Excellence (2020). *COVID-19 rapid guideline: managing the long-term effects of COVID-19.* London: NICE (https://www.nice.org.uk/guidance/ng188/resources/covid 19-rapid-guideline-managing-the-longterm-effects-ofcovid19-pdf-66142028400325, accessed 28 January 2021).

Nehme M et al. (2020). COVID-19 symptoms: Longitudinal evolution and persistence in outpatient settings. *Ann Intern Med*.M20-5926. doi: 10.7326/M20-5926.

NHS (2020a). National Guidance for post-COVID syndrome assessment clinics. London: NHS England and NHS Improvement

(https://www.england.nhs.uk/coronavirus/wpcontent/uploads/sites/52/2020/11/C0840-national-guidanc e-for-post-covid-syndrome-assesment-clinics-111220.pdf, accessed 28 January 2021).

NHS (2020b). Your COVID Recovery. London: NHS England (https://www.yourcovidrecovery.nhs.uk/your-road-to-recovery/what-next/, accessed 28 January 2021).

NIHR Leicester Biomedical Research Centre (2020). UK-REACH: UK research study into ethnicity and COVID-19 outcomes in healthcare workers. Leicester: NIHR Leicester Biomedical Research Centre (https://www.leicesterbrc.nihr.ac.uk/themes/respiratory/res

earch/uk-reach/, accessed 28 January 2021).

- Noi che il Covid lo abbiamo sconfitto (2020). SINDROME POST COVID. Facebook Group (https://www.facebook.com/groups/3044276745637072, accessed 28 January 2021).
- Office for National Statistics (2020). The prevalence of long COVID symptoms and COVID-19 complications. London: ONS

(https://www.ons.gov.uk/news/statementsandletters/thepr evalenceoflongcovidsymptomsandcovid19complications, accessed 28 January 2021).

Office for National Statistics (2021). Updated estimates of the prevalence of long COVID symptoms. London: ONS (https://www.ons.gov.uk/peoplepopulationandcommunity/ healthandsocialcare/healthandlifeexpectancies/adhocs/127 88updatedestimatesoftheprevalenceoflongcovidsymptoms , accessed 28 January 2021).

Padda I et al. (2020). The neurological complexities and prognosis of COVID-19. *SN Compr Clin Med*.1–12. doi: 10.1007/s42399-020-00527-2.

Patient led research for COVID-19 (2020). Report: What does COVID-19 recovery actually look like? (https://patientresearchcovid19.com/research/report-1/,accessed 28 January 2021).

Perego E et al. (2020). Why the patient-made term "Long Covid" is needed. *Wellcome Open Res*.5:224. doi: 10.12688/wellcomeopenres.16307.1.

Pezzini A, Padovani A (2020). Lifting the mask on neurological manifestations of COVID-19. *Nat Rev Neurol*.16(11):636–44.

PHOSP-COVID (2020). Post-hospitalisation COVID-19 study. Leicester: University of Leicester (https://www.phosp.org/, accessed 28 January 2021).

Puntmann VO et al. (2020). Outcomes of cardiovascular magnetic resonance imaging in patients recently recovered from coronavirus disease 2019 (COVID-19). JAMA Cardiol.5(11):1265–73.

Rajpal S et al. (2020). Cardiovascular magnetic resonance findings in competitive athletes recovering from COVID-19 infection. JAMA Cardiol.6(1):116–18. doi: 10.1001/jamacardio.2020.4916.

Reynolds KJ et al. (2004). The economic impact of chronic fatigue syndrome. Cost Eff Resour Alloc.2(1):4. doi: 10.1186/1478-7547-2-4.

Royal College of General Practitioners (2020). Written evidence (COV0051). London: UK Parliament (https://committees.parliament.uk/writtenevidence/12976/ html/, accessed 28 January 2021).

Royal College of Occupational Therapists (2020). *How to conserve your energy: Practical advice for people during and after having COVID-19*. London: Royal College of Occupational Therapists (https://www.rcot.co.uk/conserving-energy, accessed 28

January 2021).

Rubino F et al. (2020). New-onset diabetes in Covid-19. N Engl J Med.383(8):789–90. Sathish T et al. (2020). Proportion of newly diagnosed diabetes in COVID-19 patients: A systematic review and meta-analysis. *Diabetes Obes Metab.* doi: 10.1111/dom.14269.

SIMPLECOVID (2020). Simple COVID. London: SIMPLECOVID. (https://simplecovid.org/, accessed 28 January 2021).

Simpson F, Lokugamage A (2020). Counting long covid in children. BMJ Opin. 16 October 2020 (https://blogs.bmj.com/bmj/2020/10/16/counting-longcovid-in-children/?fbclid=IwAR0sKQOvNfHpf_LzZeswGEM qdWLwJKfSIVk03S0S6ZY6153XwS7qTj7LVZQ, accessed 28 January 2021).

Singh AK et al. (2020). Prevalence of co-morbidities and their association with mortality in patients with COVID-19: A systematic review and meta-analysis. *Diabetes Obes Metab*.22(10):1915–24.

Sivan M, Halpin SJ, Gee J (2020). Assessing long-term rehabilitation needs in covid-19 survivors using a telephone screening tool (C19-YRS tool). Adv Clin Neurosci Rehabil.19:14–17.

Sivan M, Taylor S (2020). NICE guideline on long covid. BMJ.371:m4938.

Sociedad Española de Médicos Generales y de Familia (2020a). *Kit de atención básica al paciente con enfermedad COVID-19 persistente*. Madrid: Sociedad Española de Médicos Generales y de Familia (https://www.semg.es/images/2020/Noticias/kit_at_covid_ persistente_opt.pdf, accessed 28 January 2021).

Sociedad Española de Médicos Generales y de Familia (2020b). La SEMG elabora un protocolo de atención básica para los afectados de COVID-19 persistente. Madrid: Sociedad Española de Médicos Generales y de Familia (https://www.semg.es/index.php/noticias/item/594noticia-20200929, accessed 28 January 2021).

Society of Critical Care Medicine (2020). *Post-intensive care syndrome*. Mount Prospect, IL: Society of Critical Care Medicine (https://www.sccm.org/MyICUCare/THRIVE/Post-intensive-Care-Syndrome, accessed 28 January 2021).

Sudre CH et al. (2020). Attributes and predictors of Long-COVID: analysis of COVID cases and their symptoms collected by the Covid Symptoms Study App. *medRxiv*.2020.10.19.20214494. doi: 10.1101/2020.10.19.20214494.

Swann OV et al. (2020). Clinical characteristics of children and young people admitted to hospital with covid-19 in United Kingdom: prospective multicentre observational cohort study. *BMJ*.370:m3249.

Taquet M et al. (2020). Bidirectional associations between COVID-19 and psychiatric disorder: retrospective cohort studies of 62 354 COVID-19 cases in the USA. *Lancet*.8(2):130–40. doi: S2215–0366(2220)30462– 30464.

Tenforde MW et al. (2020). Symptom duration and risk factors for delayed return to usual health among outpatients with COVID-19 in a multistate health care systems network – United States, March–June 2020. *MMWR Morb Mortal Wkly Rep.*69(30):993–8.

Townsend L et al. (2020). Persistent fatigue following SARS-CoV-2 infection is common and independent of severity of initial infection. *PLoS One*.15(11):e0240784.

- UKCDR (2020). COVID-19 Research project tracker by UKCDR & GloPID-R. London: UK Collaborative on Development Research (https://www.ukcdr.org.uk/covidcircle/covid-19-research-project-tracker/, accessed 28 January 2021).
- UKRI (2020). "Long-COVID" in our community new call for research proposals. London: UK Research and Innovation. (https://www.ukri.org/news/long-covid-in-our-communitynew-call-for-research-proposals/?fbclid=IwAR3ty4Kp2iiSVV CRsYIWZFENqpHT-
- d_WD71jn4w8FONPpj75L5G2ee2DcO8, accessed 28 January 2021).
- University of Oxford (2020). *Global consortium launches new study into long-term effects of COVID-19*. Oxford: University of Oxford. 11 September 2020. In: University of Oxford/News (https://www.ox.ac.uk/news/2020-09-11global-consortium-launches-new-study-long-term-effects-c ovid-19, accessed 28 January 2021).
- WHO (2020a). Emergency use ICD codes for COVID-19 disease outbreak. Geneva: World Health Organization (https://www.who.int/standards/classifications/classificatio n-of-diseases/emergency-use-icd-codes-for-covid-19disease-outbreak, accessed 28 January 2021).

WHO (2020b). Multisystem inflammatory syndrome in children and adolescents temporally related to COVID-19. Geneva: World Health Organization (https://www.who.int/newsroom/commentaries/detail/multisystem-inflammatory-synd rome-in-children-and-adolescents-with-covid-19, accessed 28 January 2021).

- Wong AW et al. (2020). Patient-reported outcome measures after COVID-19: a prospective cohort study. *Eur Respir* J.56(5):2003276. doi: 10.1183/13993003.03276-2020.
- Yelin D et al. (2020). Long COVID-19 it's not over until? *Clin Microbiol Infect*.S1198-743X(20)30750-3. doi: 10.1016/j.cmi.2020.12.001.

How do Policy Briefs bring the evidence together?

There is no one single way of collecting evidence to inform policymaking. Different approaches are appropriate for different policy issues, so the Observatory briefs draw on a mix of methodologies (see Figure A) and explain transparently the different methods used and how these have been combined. This allows users to understand the nature and limits of the evidence.

There are two main 'categories' of briefs that can be distinguished by method and further 'sub-sets' of briefs that can be mapped along a spectrum:

- A rapid evidence assessment: This is a targeted review of the available literature and requires authors to define key terms, set out explicit search strategies and be clear about what is excluded.
- Comparative country mapping: These use a case study approach and combine document reviews and consultation with appropriate technical and country experts. These fall into two groups depending on whether they prioritize depth or breadth.
- Introductory overview: These briefs have a different objective to the rapid evidence assessments but use a similar methodological approach. Literature is targeted and reviewed with the aim of explaining a subject to 'beginners'.

Most briefs, however, will draw upon a mix of methods and it is for this reason that a 'methods' box is included in the introduction to each brief, signalling transparently that methods are explicit, robust and replicable and showing how they are appropriate to the policy question.



Figure A: The policy brief spectrum

Source: Erica Richardson

Joint Policy Briefs

- 1. How can European health systems support investment in and the implementation of population health strategies? *David McDaid, Michael Drummond, Marc Suhrcke*
- How can the impact of health technology assessments be enhanced? Corinna Sorenson, Michael Drummond, Finn Børlum Kristensen, Reinhard Busse
- 3. Where are the patients in decision-making about their own care? Angela Coulter, Suzanne Parsons, Janet Askham
- How can the settings used to provide care to older people be balanced? Peter C. Coyte, Nick Goodwin, Audrey Laporte
- When do vertical (stand-alone) programmes have a place in health systems? *Rifat A. Atun, Sara Bennett, Antonio Duran*
- How can chronic disease management programmes operate across care settings and providers? Debbie Singh
- 7. How can the migration of health service professionals be managed so as to reduce any negative effects on supply? *James Buchan*
- 8. How can optimal skill mix be effectively implemented and why? *Ivy Lynn Bourgeault, Ellen Kuhlmann, Elena Neiterman, Sirpa Wrede*
- Do lifelong learning and revalidation ensure that physicians are fit to practise? Sherry Merkur, Philipa Mladovsky, Elias Mossialos,
- Martin McKee 10. How can health systems respond to population ageing? Bernd Rechel, Yvonne Doyle, Emily Grundy, Martin McKee
- 11. How can European states design efficient, equitable and sustainable funding systems for long-term care for older people? José-Luis Fernández, Julien Forder, Birgit Trukeschitz, Martina Rokosová, David McDaid
- 12. How can gender equity be addressed through health systems? Sarah Payne
- 13. How can telehealth help in the provision of integrated care? Karl A. Stroetmann, Lutz Kubitschke, Simon Robinson, Veli Stroetmann, Kevin Cullen, David McDaid
- 14. How to create conditions for adapting physicians' skills to new needs and lifelong learning Tanya Horsley, Jeremy Grimshaw, Craig Campbell
- How to create an attractive and supportive working environment for health professionals *Christiane Wiskow, Tit Albreht, Carlo de Pietro*
- How can knowledge brokering be better supported across European health systems?
 John N. Lavis, Govin Permanand, Cristina Catallo, BRIDGE Study Team
- 17. How can knowledge brokering be advanced in a country's health system? John. N Lavis, Govin Permanand, Cristina Catallo, BRIDGE Study Team
- How can countries address the efficiency and equity implications of health professional mobility in Europe? Adapting policies in the context of the WHO Code and EU freedom of movement *Irene A. Glinos, Matthias Wismar, James Buchan, Ivo Rakovac*
- 19. Investing in health literacy: What do we know about the co-benefits to the education sector of actions targeted at children and young people? David McDaid
- 20. How can structured cooperation between countries address health workforce challenges related to highly specialized health care? Improving access to services through voluntary cooperation in the EU.
 - Marieke Kroezen, James Buchan, Gilles Dussault, Irene Glinos, Matthias Wismar
- 21. How can voluntary cross-border collaboration in public procurement improve access to health technologies in Europe? Jaime Espín, Joan Rovira, Antoinette Calleja, Natasha Azzopardi-Muscat, Erica Richardson, Willy Palm, Dimitra Panteli

- 22. How to strengthen patient-centredness in caring for people with multimorbidity in Europe? Iris van der Heide, Sanne P Snoeijs, Wienke GW Boerma, François GW Schellevis, Mieke P Rijken. On behalf of the ICARE4EU consortium
- 23. How to improve care for people with multimorbidity in Europe? Mieke Rijken, Verena Struckmann, Iris van der Heide, Anneli Hujala, Francesco Barbabella, Ewout van Ginneken, François Schellevis. On behalf of the ICARE4EU consortium
- 24. How to strengthen financing mechanisms to promote care for people with multimorbidity in Europe? Verena Struckmann, Wilm Quentin, Reinhard Busse, Ewout van Ginneken. On behalf of the ICARE4EU consortium
- 25. How can eHealth improve care for people with multimorbidity in Europe? Francesco Barbabella, Maria Gabriella Melchiorre, Sabrina
 - Quattrini, Roberta Papa, Giovanni Lamura. On behalf of the ICARE4EU consortium
- 26. How to support integration to promote care for people with multimorbidity in Europe? Anneli Hujala, Helena Taskinen, Sari Rissanen. On behalf of the ICARE4EU consortium
- 27. How to make sense of health system efficiency comparisons? Jonathan Cylus, Irene Papanicolas, Peter C Smith
- 28. What is the experience of decentralized hospital governance in Europe?
- Bernd Rechel, Antonio Duran, Richard Saltman
 29. Ensuring access to medicines: How to stimulate innovation to meet patients' needs? Dimitra Panteli, Suzanne Edwards
- 30. Ensuring access to medicines: How to redesign pricing, reimbursement and procurement? Sabine Vogler, Valérie Paris, Dimitra Panteli
- 31. Connecting food systems for co-benefits: How can food systems combine diet-related health with environmental and economic policy goals? *Kelly Parsons, Corinna Hawkes*
- 32. Averting the AMR crisis: What are the avenues for policy action for countries in Europe? Michael Anderson, Charles Clift, Kai Schulze, Anna Sagan, Saskia Nahrgang, Driss Ait Ouakrim, Elias Mossialos
- 33. It's the governance, stupid! TAPIC: a governance framework to strengthen decision making and implementation Scott L. Greer, Nikolai Vasev, Holly Jarman, Matthias Wismar, Josep Figueras
- 34. How to enhance the integration of primary care and public health? Approaches, facilitating factors and policy options *Bernd Rechel*
- 35. Screening. When is it appropriate and how can we get it right? Anna Sagan, David McDaid, Selina Rajan, Jill Farrington, Martin McKee
- Strengthening health systems resilience: key concepts and strategies Steve Thomas, Anna Sagan, James Larkin, Jonathan Cylus, Josep Figueras, Marina Karanikolos
- 37. Building on value-based health care Peter C Smith, Anna Sagan, Luigi Siciliani, Dimitra Panteli, Martin McKee, Agnès Soucat, Josep Figueras
- Regulating the unknown: A guide to regulating genomics for health policy-makers
 Gemma A Williams, Sandra Liede, Nick Fahy, Kristiina Aittomaki, Markus Perola

The European Observatory has an independent programme of policy briefs and summaries which are available here: http://www.euro.who.int/en/about-us/partners/ observatory/publications/policy-briefs-and-summaries World Health Organization Regional Office for Europe UN City, Marmorvej 51, DK-2100 Copenhagen Ř, Denmark Tel.: +45 39 17 17 17 Fax: +45 39 17 18 18 E-mail: postmaster@euro.who.int web site: www.euro.who.int

The European Observatory on Health Systems and Policies is a partnership that supports and promotes evidence-based health policy-making through comprehensive and rigorous analysis of health systems in the European Region. It brings together a wide range of policy-makers, academics and practitioners to analyse trends in health reform, drawing on experience from across Europe to illuminate policy issues. The Observatory's products are available on its web site (http://www.healthobservatory.eu).



Web ISSN 1997-8073

INQ000249018_0030