

- wheezing;
- swelling of your lips, face, tongue or throat;
- hives or rash;
- nausea or vomiting;
- stomach pain.

Talk to your doctor or nurse if you develop any other side effects. These can include:

**Very common** (may affect more than 1 in 10 people):

- swelling/tenderness of the underarm glands
- decreased appetite (observed in 6 month to 5 year olds)
- irritability/crying (observed in 6 month to 5 year olds)
- headache
- sleepiness (observed in 6 month to 5 year olds)
- nausea
- vomiting
- muscle ache, joint aches, and stiffness
- pain or swelling at the injection site
- redness at the injection site (some of which may occur approximately 9 to 11 days after the injection)
- feeling very tired
- chills
- fever

**Common** (may affect up to 1 in 10 people):

- diarrhoea
- rash
- rash or hives at the injection site (some of which may occur approximately 9 to 11 days after the injection)

**Uncommon** (may affect up to 1 in 100 people):

- itchiness at the injection site
- dizziness
- stomach pain
- raised, itchy rash (urticaria) (which may occur from the time of injection and up to approximately two weeks after the injection)

**Rare** (may affect up to 1 in 1,000 people)

- temporary one-sided facial drooping (Bell's palsy)
- swelling of the face (Swelling of the face may occur in patients who have had facial cosmetic injections.)
- decreased sense of touch or sensation
- unusual feeling in the skin, such as tingling or a crawling feeling (paraesthesia)

**Very rare** (may affect up to 1 in 10,000 people)

- inflammation of the heart muscle (myocarditis) or inflammation of the lining outside the heart (pericarditis) which can result in breathlessness, palpitations or chest pain

**Frequency not known**

- severe allergic reactions with breathing difficulties (anaphylaxis)
- reaction of increased sensitivity or intolerance by the immune system (hypersensitivity)
- a skin reaction that causes red spots or patches on the skin that may look like a target or "bulls-eye" with a dark red centre surrounded by paler red rings (erythema multiforme)
- heavy menstrual bleeding (most cases appeared to be non-serious and temporary in nature)
- extensive swelling of the vaccinated limb